arXiv:1211.4520v1 [cs.NE] 19 Nov 2012

Storing Cycles in Hopfield-type Networks with Pseudoinverse Learning Rule: Admissibility and Network Topology

Chuan Zhang[a], Gerhard Dangelmayr[a], Iuliana Oprea[a,*]

[a]*Department of Mathematics, Colorado State University, Fort Collins, CO 80523, USA*

Abstract

The storage of binary cycles in continuous-time Hopfield-type networks is investigated. We call a cycle defined by a binary matrix Σ admissible if a connectivity matrix satisfying the cycle's transition conditions exists, and if so construct it using the pseudoinverse learning rule. Our main focus is on the structural features of admissible cycles and the topology of the corresponding networks. We show that Σ is admissible if and only if its discrete Fourier transform contains exactly $r = \text{rank}(\Sigma)$ nonzero columns. Based on the decomposition of the rows of Σ into disjoint subsets corresponding to loops, where a loop is defined by the set of all cyclic permutations of a row, cycles are classified as simple cycles, and separable or inseparable composite cycles. Simple cycles contain rows from one loop only, and the network topology is a feedforward chain with feedback to one neuron if the loop-vectors in Σ are cyclic permutations of each other. For special cases this topology simplifies to a ring with only one feedback. Composite cycles contain rows from at least two disjoint loops, and the neurons corresponding to the loop-vectors in Σ from the same loop are identified with a cluster. Networks constructed from separable composite cycles decompose into completely isolated clusters. For inseparable composite cycles at least two clusters are connected, and the cluster-connectivity is related to the intersections of the spaces spanned by the loop-vectors of the clusters.

Keywords: Cyclic Patterns, Hopfield-type Networks, Pseudoinverse Learning Rule, Admissibility, Network Topology

---

[*] Corresponding author
Email address: `juliana@math.colostate.edu` (Iuliana Oprea)

---

1. Introduction

Applications of artificial neural networks in content addressable (associative) memory have attracted much attention in the last few decades (Hopfield, 1982, 1984; Little, 1974; McEliece et al., 1987; Šíma & Orponen, 2003; López-Rodríguez et al., 2005). Hopfield-type networks are among the most popular models of artificial neural networks for studying content addressable memory. According to Hopfield's original idea, the privileged regime to store information has been fixed point attractors, however experiments (e.g., Korn & Faure, 2003) indicate that cycles are used to store information and chaotic dynamics appears as the background regime composed of these cyclic "memory bags".

In both discrete and continuous asymmetric variants of Hopfield networks, the storage and retrieval of complex sequences including cycles of binary patterns have been investigated (Personnaz et al., 1986; Guyon et al., 1988; Gencic et al., 1990), and biologically plausible learning rules such as Hebb's rule, the pseudoinverse rule and their variants with or without delays have been used. While simulations of networks constructed using Hebbian learning rules have been shown to be qualitatively consistent with experimental recordings (e.g. Kleinfeld, 1986; Kleinfeld & Sompolinsky, 1988), it is well known (e.g. Rojas, 1996) that Hopfield networks with Hebbian rules do not perform well when the patterns to be stored are correlated which is usually the case in practice. Moreover, it has been recently suggested that the pseudoinverse learning rule and its variants may take key roles in the associative perception of human faces in the human cortex (Zifan et al., 2007) and the encoding of location information in the rat hippocampus (Marinaro et al., 2007). Therefore, in this and a subsequent paper (Zhang et al., 2013), we study the storage and retrieval of cycles in continuous-time Hopfield-type networks using the pseudoinverse learning rule.

Although the pseudoinverse rule and its variants (Kohonen, 1988; Personnaz et al., 1986; Diederich & Opper, 1987) extend to more general cases, most investigations in discrete-time Hopfield-type networks characterized or were implemented for cycles or sequences of linearly independent patterns (e.g. Personnaz et al., 1986; Kanter & Sompolinsky, 1987; Diederich & Opper, 1987). An approach to storing cycles of correlated as well as linearly independent patterns in continuous-time Hopfield-type networks has been

proposed by Gencic et al. (1990). In this study, a successfully retrieved cycle is revealed as an attracting limit cycle in the network dynamics, but it remains unclear under which conditions the storage and retrieval of a cycle is possible. On the one hand, a connectivity matrix in accordance with the cycle's transition conditions may not exist, and on the other hand, as shown by Zhang et al. (2013), there are cycles of linearly independent patterns for which a connectivity matrix can be constructed, however, the dynamics of the corresponding networks without delay do not show an attracting limit cycle. Accordingly, understanding the storage and retrieval of cycles in continuous-time Hopfield-type networks requires to address the following questions:
(1) For a given cycle, does there exist a connectivity matrix for a Hopfield-type network in accordance with the cycle's transition conditions? If the answer to this question is affirmative, we call the cycle admissible and construct its connectivity matrix using the pseudoinverse learning rule.
(2) If a cycle is admissible, is it indeed retrieved successfully in the corresponding continuous-type Hopfield-type network, that is, does the network dynamics exhibit an attracting limit cycle that resembles the prescribed transitions between the cycle's patterns?

We address these questions in this and a subsequent paper (Zhang et al., 2013). The focus of the present paper is on the first question. We formulate and prove sufficient and necessary conditions for admissibility of cycles, and study the topology of the corresponding networks. In the sequel to this paper, the bifurcations in continuous-time Hopfield-type networks with and without delay are analyzed, which reveals how oscillations corresponding to the transitions prescribed by the cycle are created, and which cycles can be retrieved successfully. Our main findings are (i) successfully retrieved cycles are created in a Hopf bifurcation from the origin, (ii) for networks with ring-topology they are destroyed in a saddle-node on limit cycle bifurcation, and (iii) most admissible cycles are retrieved in networks with delays for sufficiently large delay-time, even if they are not retrieved in networks without delay.

In our study, a cycle is defined by a matrix, Σ, of binary states for the firing rates. To store the cycle in a continuous-time Hopfield-type network, the connectivity matrix is decomposed into a weighted sum of two components. The first component serves to store the individual states in Σ as fixed points, which is implemented using the standard pseudoinverse rule. The second, cycle-component serves to implement the transition conditions, where we stipulate that the states in Σ are traversed in a cyclic manner. These conditions lead to a matrix-equation, which has a solution if and only if the rows of Σ span the same space as the cyclic permutations of the rows of Σ. If this condition is satisfied, Σ is admissible, and we construct a solution of the matrix-equation for the cycle-component of the connectivity matrix using again the pseudoinverse method. For the discrete-time networks studied by Personnaz et al. (1986) and Guyon et al. (1988) only the cycle-component is needed to store a cycle, however in the continuous case the fixed point component is needed as well.

Our main objective is to study the structural features of admissible cycles Σ and the topology of the networks constructed from them. Since the storage of cycles in the discrete-time networks considered by Personnaz et al. (1986) and Guyon et al. (1988) leads to the same equation for the connectivity matrix as for the cycle-component in the continuous case, our results are also applicable to discrete networks. A basic result is that if and only if the discrete Fourier transform of Σ contains exactly $r = \text{rank}(\Sigma)$ nonzero columns, then Σ is admissible.

Our approach to classify cycles is based on the decomposition of the row vectors of Σ into disjoint subsets corresponding to different loops created by cyclic permutations of the rows. If the cycle is admissible, each of these loops is associated with an invariant subspace of the row space, $\mathcal{U}$, of Σ under cyclic permutations. This row-decomposition leads naturally to a classification of cycles into simple cycles, separable composite cycles, and inseparable composite cycles. Simple cycles contain rows from a single loop only. Composite cycles contain rows from at least two disjoint loops, and for each loop the neurons corresponding to the loop vectors in Σ are identified with a cluster, which in turn corresponds to an indecomposable invariant subspace of $\mathcal{U}$ under cyclic permutations if Σ is admissible. Two clusters are directly connected if their subspaces intersect nontrivially, and they are connected if they are part of a chain of directly connected clusters. A network constructed from a simple admissible cycle has only one cluster, and we show that the network topology is a feedforward chain with feedback to one neuron if the loop vectors in Σ are all cyclic permutations of each other. For special simple cycles, there is only one feedback and the network topology simplifies to a ring. Networks constructed from separable composite cycles decompose into completely isolated clusters.

The paper is organized as follows. In Section 2, the equations for the networks and the pseudoinverse learning rule are introduced. In Section 3, the general admissibility criterion in terms of the discrete Fourier transform of the cycle matrix is formulated and proved, and the relation of admissible cycles with cyclic permutation groups is discussed. Based on the structural features of the invariant subspaces of the row space of an admissible cycle, in Section 4, admissible cycles are classified into simple cycles, and separable and inseparable composite cycles, and for each type of cycles a corresponding admissibility condition is derived. In Section 5, we study the topology of networks constructed from different types of admissible cycles, and in Section 6 we discuss some implications of the results presented in this paper.

2. Hopfield-type Networks and the Pseudoinverse Learning Rule

A continuous-time Hopfield-type network (Hopfield, 1984) is described by a system of ordinary differential equations for $u_i(t)$, $1 \le i \le N$, which model the membrane potential of the i-th neuron in the network at time t:

$$C_i \frac{du_i}{dt} = -\frac{u_i}{R_i} + \sum_{j=1}^{N} \tilde{J}_{ij} v_j + I_i,\ i = 1, 2, \ldots, N, \tag{1}$$

where $N \ge 2$ is the number of neurons in the network, $C_i > 0$ and $R_i > 0$ are the neuron amplifier input capacitance and resistance of the i-th neuron respectively, $v_j(t)$ models the firing rate of the j-th neuron, and I_i is an external input fed into the i-th neuron. The connectivity matrix $\tilde{\mathbf{J}} = (\tilde{J}_{ij})_{N\times N}$ describes the connection strengths of the network. The sign of $\tilde{J}_{ij}$ characterizes the polarity of the synaptic connection from the j-th neuron to the i-th neuron. That is, if the output from the j-th neuron excites the i-th neuron, then $\tilde{J}_{ij} > 0$, and if it inhibits, then $\tilde{J}_{ij} < 0$.

The firing rate $v_j(t)$ is related to the membrane potential $u_j(t)$ through a sigmoid-shaped gain function, $v_j = g(u_j)$. Following Hopfield's original smoothness assumption, we choose $g(u) = \tanh(\lambda u)$, where the parameter λ controls the steepness of the sigmoid function. As further simplification to (1) we normalize $R_i = C_i = 1$ and neglect external inputs, i.e. $I_i = 0$ for every i.

With these simplifications, (1) becomes (dots denote time derivatives)

$$\dot{u}_i = -u_i + \sum_{j=1}^{N} \tilde{J}_{ij} \tanh(\lambda u_j). \tag{2}$$

Alternatively, since $u_j = \text{arctanh}(v_j)/\lambda$, (2) can be rewritten as a system of differential equations for the firing rates,

$$\dot{v}_i = \lambda(1 - v_i^2)\left(\sum_{j=1}^{N} \tilde{J}_{ij} v_j - \frac{1}{\lambda}\text{arctanh}(v_i)\right). \tag{3}$$

Using vector notation, $\mathbf{u} = (u_1, \ldots, u_N)^T$, $\mathbf{v} = (v_1, \ldots, v_N)^T$, (2) can be more compactly written as

$$\dot{\mathbf{u}} = -\mathbf{u} + \tilde{\mathbf{J}} \tanh(\lambda \mathbf{u}), \tag{4}$$

where here and subsequently a scalar function of a vector or matrix denotes the vector or matrix obtained by applying the function to each component, i.e.

$$\tanh(\lambda \mathbf{u}) = (\tanh(\lambda u_1), \ldots, \tanh(\lambda u_N))^T.$$

In this paper, we study the structure of binary pattern cycles that can be stored in the network modeled by the above autonomous system. Following Hopfield (1982, 1984), any N-dimensional $\{-1, 1\}$-valued column vector is identified with a *binary vector* or *pattern*, and we use $+$ and $-$ to denote 1 and -1. A *sequence* of q binary patterns is defined by $q-1$ consecutive transitions, $\xi^{(1)} \to \xi^{(2)} \to \cdots \to \xi^{(q)}$, with $\xi^{(\mu)} = (\xi_1^{(\mu)}, \xi_2^{(\mu)}, \ldots, \xi_N^{(\mu)})^T$, $\xi_i^{(\mu)} = +$ or $-$, and $\mu = 1, 2, \ldots, q$. A matrix pair (Σ, F) is called the *matrix form* of a sequence, if $\Sigma = (\xi^{(1)}, \xi^{(2)}, \ldots, \xi^{(p)})$, $F = (f^{(1)}, f^{(2)}, \ldots, f^{(p)})$ and the two matrices are related through the transition conditions, $\xi^{(\mu)} \to f^{(\mu)}$, $\mu = 1, 2, \ldots, p$. In this formulation multiple sequences can be combined, and if terminal states are present one has to include the condition $\xi^{(q)} \to \xi^{(q)}$, so that for each μ a transition $\xi^{(\mu)} \to f^{(\mu)} = \xi^{(\nu(\mu))}$ exists.

According to Personnaz et al. (1986), the transition conditions can be conveniently formulated in terms of a $p \times p$ transition matrix $\mathbf{P}$ as $F = \Sigma\mathbf{P}$, where $\mathbf{P}_{\nu\mu} = 1$ if $f^{(\mu)} = \xi^{(\nu)}$ and 0 otherwise. Personnaz et al. (1986) and Guyon et al. (1988) have studied the storage of sequences in discrete-time binary networks. In these papers, the transition conditions have been reduced to the single matrix equation

$$\mathbf{J}\Sigma = F. \tag{5}$$

It was pointed out by Personnaz et al. (1986), that, if $F\Sigma^+\Sigma = F$, where Σ^+ is the Moore-Penrose pseudoinverse of Σ, then (5) has the exact solution $\tilde{\mathbf{J}} = F\Sigma^+$, which was called *associating learning rule* by these authors.

A sequence is called a *cycle* of period p, if there exists a smallest positive integer p such that for every $\mu \in \mathbb{N}$, $\xi^{(\mu+p)} = \xi^{(\mu)}$. Accordingly, a cycle



Figure 1: A successfully retrieved cycle Σ (see text for details) in a network of three neurons. **A** shows the firing rates $v_i(t)$ of the neurons and the raster plot **B** the overlaps, which measure the similarity of the network state $\mathbf{v}(t)$ with each of the six patterns in the cycle. **C** illustrates the retrieval of the cycle in the phase space of the system.

of period p has the matrix form (Σ, F), where $\Sigma = (\xi^{(1)}, \xi^{(2)}, \cdots, \xi^{(p)})$ and $F = (\xi^{(2)}, \xi^{(3)}, \cdots, \xi^{(p)}, \xi^{(1)})$, and the corresponding transition matrix is

$$\mathbf{P} = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{pmatrix}. \tag{6}$$

We are interested in the storage of cycles in the continuous-time Hopfield networks defined by (4). Our approach to compute a connectivity matrix $\tilde{\mathbf{J}}$ for this purpose follows Gencic et al. (1990). In this paper, the connectivity matrix is decomposed as

$$\tilde{\mathbf{J}} = \beta_K (C_0 \mathbf{J}^\mathbf{0} + C_1 \mathbf{J}), \tag{7}$$

where $\mathbf{J}^\mathbf{0}$ serves to stabilize the network in its current memory state and $\mathbf{J}$ imposes the transitions between the memory states. Here, $C_1 = 1 - C_0$ and C_0, $0 \leq C_0 \leq 1$, control the relative contributions of the two components of $\tilde{\mathbf{J}}$. The fixed point condition is realized by requiring that $\mathbf{v} = \beta_1 \xi^{(\mu)}$, with a parameter $0 < \beta_1 < 1$, is a fixed point if $C_0 = 1$. Noting that $\operatorname{arctanh}(x)$ is an odd function and $|\xi_i^{(\mu)}| = 1$, this leads, according to (3), to the condition

$$\tilde{\mathbf{J}}\beta_1\xi^{(\mu)} = \frac{1}{\lambda}\text{arctanh}(\beta_1\xi^{(\mu)}) = \frac{1}{\lambda}\text{arctanh}(\beta_1)\xi^{(\mu)},$$

for every μ, hence $\tilde{\mathbf{J}}\Sigma = \beta_K\Sigma$ with $\beta_K = \dfrac{1}{\lambda\beta_1}\text{arctanh}(\beta_1)$, which has the solution $\tilde{\mathbf{J}} = \beta_K\mathbf{J^0}$ with

$$\mathbf{J^0} = \Sigma\Sigma^+. \tag{8}$$

Regarding the transition condition, we stipulate that $\mathbf{v}(t) = \beta_1\xi^{(\mu)}$ implies $\mathbf{v}(t+\tau) = \beta_1 f^{(\mu)}$ for some τ, and require accordingly for $C_0 = 0$ that $\tilde{\mathbf{J}}\Sigma = \beta_K F$. This leads to equation (5) with $\tilde{\mathbf{J}} = \mathbf{J}$, which in terms of the transition matrix $\mathbf{P}$, equation (6), becomes

$$\mathbf{J}\Sigma = \Sigma\mathbf{P}. \tag{9}$$

According to the associating learning rule of Personnaz et al. (1986), (9) has the solution

$$\mathbf{J} = \Sigma\mathbf{P}\Sigma^+, \tag{10}$$

provided that $\Sigma\mathbf{P}\Sigma^+\Sigma = \Sigma\mathbf{P}$. If this condition is not satisfied, (9) has no solution.

The main objective of this paper is the study of the existence and properties of the solutions of (9) along with the structural features of the corresponding cycles, and the network topologies resulting from $\mathbf{J^0}$ and $\mathbf{J}$ defined by (8) and (10). A bifurcation analysis and a study of the dynamics of (4) with (7), with C_0 and $\beta = \lambda\beta_K > 1$ treated as parameters, are given in Zhang et al. (2013). In this paper also the extension of (4) to a dynamical system with a delay,

$$\dot{\mathbf{u}} = -\mathbf{u} + C_0\beta_K\mathbf{J^0}\tanh(\lambda\mathbf{u}) + C_1\beta_K\mathbf{J}\tanh(\lambda\mathbf{u}_\tau), \tag{11}$$

where $\tau > 0$ is the delay-time and $\mathbf{u}_\tau(t) = \mathbf{u}(t-\tau)$, is considered. Here we show only one example of a successfully retrieved cycle for (4) and devote the remainder of the paper to a mathematical analysis of equation (9). The example is a network of $N = 3$ neurons and the cycle consists of six states, $\Sigma = (\xi^{(1)}, \ldots, \xi^{(6)})$, with $\xi^{(1)} = (+,+,+)^T$, $\xi^{(2)} = (+,+,-)^T$, $\xi^{(3)} = (+,-,-)^T$ and $\xi^{(3+\mu)} = -\xi^{(\mu)}$ for $\mu = 1,2,3$. The retrieval of the cycle is illustrated in Figure 1, where in general the overlap of the network with the ν-th pattern is defined as $m^{(\nu)}(t) = (\xi^{(\nu)})^T\mathbf{v}(t)/N$ (Zhang et al., 2013). The raster plot of the overlaps in Figure 1B as well as the time series in Figure 1A clearly illustrate that the cycle is retrieved successfully. The parameters C_0 and β used in this simulation were $C_0 = 0.6$ and $\beta = 4$.

In the next section, we derive conditions under which a network for storing a given prescribed cycle, i.e. a connectivity matrix $\mathbf{J}$ satisfying (9) can be constructed, and discuss the structure of the irreducible subspaces of the row and column spaces of these cycles under the action of cyclic permutation groups.

3. Admissible Cycles and Cyclic Permutation Groups

Definition 1. Let $\mathbf{P}$ be the cyclic $p \times p$-permutation matrix defined in (6). A cycle defined by a binary $N \times p$-matrix $\Sigma = (\xi^{(1)}, \xi^{(2)}, \ldots, \xi^{(p)})$ is said to be *admissible*, if there is a real $N \times N$ matrix $\mathbf{J}$ such that equation (9) is satisfied.

Note that if Σ is admissible, the solution to (9) may be not unique. If there are several solutions, we select (10) as distinguished solution because of its close relationship to $\mathbf{J^0}$, see Remark 1(b) below.

Gencic et al. (1990) consider a special type of cycles defined by p binary vectors $\Sigma' = (\xi^{(1)}, \xi^{(2)}, \ldots, \xi^{(p)})$, which satisfy the transition condition $\xi^{(1)} \to \xi^{(2)} \to \cdots \xi^{(p)} \to -\xi^{(1)} \to -\xi^{(2)} \to \cdots - \xi^{(p)} \to \xi^{(1)}$. In this paper we consider these cycles as special cases of cycles of period $2p$ with $\Sigma = (\Sigma', -\Sigma')$.

For storing sequences, Personnaz et al. (1986) pointed out that, if the associating learning rule $F\Sigma^+\Sigma = F$ is satisfied, the rows of F are linear combinations of the rows of Σ. This follows from the fact that $\Sigma^+\Sigma$ is the orthogonal projection matrix onto the subspace of $\mathbb{R}^p$ spanned by the rows of Σ. For storing single cycles, their conclusion can be reformulated geometrically as follows:

Proposition 1. *A cycle Σ of size $N \times p$ is admissible, if and only if its row space is invariant under $\mathbf{P}$, i.e.*

$$\operatorname{span}\{\mathrm{R}(\Sigma)\} = \operatorname{span}\{\mathrm{R}(\Sigma\mathbf{P})\}, \tag{12}$$

where $\mathrm{R}(\Sigma)$ *denotes the set of all row vectors of* Σ.

Note that $|\mathrm{R}(\Sigma)| \leq N$, and if $|\mathrm{R}(\Sigma)| < N$ then two or more different rows of Σ are identical, that is, the corresponding neurons traverse the same cycle. Although this is a kind of redundancy, we do not exclude this possibility in our general discussion.

We next formulate a useful alternative admissibility criterion involving the eigenspaces of $\mathbf{P}$. Since $\mathbf{P}$ is a circulant matrix, $\mathbf{P}$ has the orthogonal eigenvectors $v^{(k)} = (1, \rho^k, \rho^{2k}, \ldots, \rho^{(p-1)k})^T$ for $0 \leq k < p$, where $\rho = e^{2\pi \mathbf{i}/p}$ ($\mathbf{i} = \sqrt{-1}$) is the basic primitive p-th root of unity (Lancaster & Tismenetsky, 1985), and in this special case the eigenvalues are $\bar{\rho}^k$. We set $V = (v^{(0)}, v^{(1)}, \ldots, v^{(p-1)})$, $\lambda_j = \bar{\rho}^{j-1}$ $(1 \leq j \leq p)$, $\Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_p)$ and note that $\mathbf{P} = V\Lambda V^{-1}$ with $V^{-1} = V^*/p$, where here and subsequently complex conjugation is marked by an overbar and an asterisk denotes the adjoint (complex conjugate transpose) matrix or vector.

Since the transition matrix $\mathbf{P}$ leaves its eigenspaces invariant, it follows that if the row space of Σ coincides with the direct sum of its projections onto the eigenspaces of $\mathbf{P}$, then Σ is admissible. Based on this consideration, we obtain the following admissibility criterion.

Theorem 1. *Let Σ be a cycle whose matrix form is of size $N \times p$, and let $\hat{\Sigma} = \Sigma V$. Then Σ is admissible, if and only if $\hat{\Sigma}$ has precisely r nonzero columns, where $r = \operatorname{rank}(\Sigma) = \operatorname{rank}(\hat{\Sigma})$.*

Proof. Noting that $\mathbf{J}\Sigma = \Sigma\mathbf{P} = \Sigma V\Lambda V^{-1}$ implies $\mathbf{J}\hat{\Sigma} = \hat{\Sigma}\Lambda$, it follows that Σ is admissible if and only if there exists an $N \times N$-matrix $\mathbf{J}$ such that

$$\mathbf{J}\hat{\Sigma} = \hat{\Sigma}\Lambda, \tag{13}$$

which implies

$$\mathbf{J}\mathrm{col}_j(\hat{\Sigma}) = \lambda_j \mathrm{col}_j(\hat{\Sigma}),\ 1 \leq j \leq p, \tag{14}$$

where $\mathrm{col}_j(\hat{\Sigma})$ denotes the j-th column of $\hat{\Sigma}$.

Suppose now that Σ is admissible and $\operatorname{rank}(\Sigma) = r$. Since the columns of V consist of eigenvectors associated to distinct eigenvalues of $\mathbf{P}$, it follows that $\hat{\Sigma} = \Sigma V$ has r linearly independent columns. Assume $\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma})$ are linearly independent. Then (14) implies that $\lambda_{\mu_1}, \ldots, \lambda_{\mu_r}$ are eigenvalues of $\mathbf{J}$ and $\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma})$ are the corresponding eigenvectors. If $\hat{\Sigma}$ has an additional nonzero column, $\mathrm{col}_j(\hat{\Sigma})$, with $j \notin \{\mu_1, \ldots, \mu_r\}$, then by (14) this column is an eigenvector of $\mathbf{J}$ corresponding to the eigenvalue λ_j and $\lambda_j \neq \lambda_{\mu_i}$ for $1 \leq i \leq r$. On the other hand, $\mathrm{col}_j(\hat{\Sigma})$ is a linear combination of $\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma})$ which is impossible, since eigenvectors corresponding to different eigenvalues are linearly independent. Thus all columns except $\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma})$ must be zero.

Conversely, assume $\hat{\Sigma}$ has r nonzero columns $\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma})$ and all other columns of $\hat{\Sigma}$ are zero. Since $\operatorname{rank}(\hat{\Sigma}) = r$, these columns are linearly independent. Let Q be a $p \times p$ permutation matrix that maps the column j to the column μ_j for $1 \leq j \leq r$. Then

$$\begin{aligned}\hat{\Sigma} &= [0, \ldots, 0, \mathrm{col}_{\mu_1}(\hat{\Sigma}), 0, \ldots, 0, \mathrm{col}_{\mu_r}(\hat{\Sigma}), 0, \ldots, 0] \\ &= [\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma}), 0, \ldots, 0]Q.\end{aligned}$$

Let $\hat{\Sigma}_0 = [\mathrm{col}_{\mu_1}(\hat{\Sigma}), \ldots, \mathrm{col}_{\mu_r}(\hat{\Sigma})]$ and define

$$\mathbf{J} = \hat{\Sigma}\Lambda Q^T \begin{bmatrix} (\hat{\Sigma}_0^*\hat{\Sigma}_0)^{-1}\hat{\Sigma}_0^* \\ O_{(p-r)\times N} \end{bmatrix}.$$

Then

$$\begin{aligned}\mathbf{J}\hat{\Sigma} &= \hat{\Sigma}\Lambda Q^T \begin{bmatrix} (\hat{\Sigma}_0^*\hat{\Sigma}_0)^{-1}\hat{\Sigma}_0^* \\ O_{(p-r)\times N} \end{bmatrix} [\hat{\Sigma}_0, O_{N\times(p-r)}]Q \\ &= \hat{\Sigma}\Lambda Q^T \begin{bmatrix} I_{r\times r} & O_{r\times(p-r)} \\ O_{(p-r)\times r} & O_{(p-r)\times(p-r)} \end{bmatrix} Q \\ &= \hat{\Sigma}\Lambda \mathrm{diag}(s_1, \ldots, s_p) \\ &= \hat{\Sigma}\Lambda.\end{aligned}$$

where $I_{r\times r}$ is the identity matrix of size $r \times r$, $O_{m\times n}$ is zero matrix of size $m \times n$, and

$$s_j = \begin{cases} 1, \mathrm{if} j \in \{\mu_1, \ldots, \mu_r\} \\ 0, \mathrm{if} j \notin \{\mu_1, \ldots, \mu_r\} \end{cases}.$$

□

Remark 1. A group theoretical interpretation of admissible cycles Σ and the associated matrices $\mathbf{J^0}$ and $\mathbf{J}$ can be given as follows:
(a) We denote by $\mathbb{Z}_p$ the cyclic group of order p defined by addition of integers modulo p. Viewed as permutation group, the generator of $\mathbb{Z}_p$, addition by 1 mod p, corresponds to the cyclic permutation $\{0, 1, \ldots, p-1\} \rightarrow \{1, 2, \ldots, p-1, 0\}$, and the matrix $\mathbf{P}$ is an orthogonal representation of this generator in $\mathbb{R}^p$ or $\mathbb{C}^p$ that cyclically permutes row vectors to the left. Accordingly, the matrices $\mathbf{P}^k$, $0 \leq k < p$, form a p-dimensional representation of $\mathbb{Z}_p$ with $\mathbf{P}^p = I$, the $p \times p$ identity matrix.
(b) The admissibility condition $\mathrm{span}\{\mathrm{R}(\Sigma)\} = \mathrm{span}\{\mathrm{R}(\Sigma\mathbf{P})\}$ means that the rows of Σ span a subspace of $\mathbb{R}^p$ or $\mathbb{C}^p$ that is invariant under $\mathbf{P}$, and hence under the full representation of $\mathbb{Z}_p$, that is,

$$\mathrm{span}\{\mathrm{R}(\Sigma)\} = \mathrm{span}\{\mathrm{R}(\Sigma\mathbf{P}^k)\},\ 0 \leq k < p.$$

Moreover, with $\mathbf{J} = \Sigma\mathbf{P}\Sigma^+$ and $\Sigma\mathbf{P}\Sigma^+\Sigma = \Sigma\mathbf{P}$, we find that $\mathbf{J}^2 = \Sigma\mathbf{P}^2\Sigma^+$ and inductively

$$\mathbf{J}^k = \Sigma\mathbf{P}^k\Sigma^+,\ 0 \leq k < p, \tag{15}$$

which shows that $\mathbf{J}^p = \Sigma\Sigma^+ = \mathbf{J}^{\mathbf{0}}$ and $\mathbf{J}^k\Sigma = \Sigma\mathbf{P}^k$ for $0 \leq k < p$. This means that $\mathbf{J}$ restricted to the column space span$\{\mathrm{C}(\Sigma)\}$, where $\mathrm{C}(\Sigma)$ denote the set of column vectors of Σ, generates a representation of $\mathbb{Z}_p$ in this subspace of $\mathbb{R}^N$. We also note that $\mathbf{J}^{\mathbf{0}}$ is the orthogonal projection onto span$\{\mathrm{C}(\Sigma)\}$, and

$$\mathbf{J}^{\mathbf{0}}\mathbf{J}^k = \mathbf{J}^k\mathbf{J}^{\mathbf{0}} = \mathbf{J}^k, \tag{16}$$

for all $0 \leq k < p$, which is a straightforward consequence of the basic properties $\Sigma\Sigma^+\Sigma = \Sigma$ and $\Sigma^+\Sigma\Sigma^+ = \Sigma^+$ of the pseudoinverse and (15). Clearly, the ranks of $\mathbf{J}$, $\mathbf{J}^{\mathbf{0}}$ and Σ coincide and are equal to the dimensions of the vector spaces span$\{\mathrm{C}(\Sigma)\}$ and span$\{\mathrm{R}(\Sigma)\}$ in which $\mathbb{Z}_p$ acts with matrix generators $\mathbf{J}$ and $\mathbf{P}$, respectively.
(c) The group $\mathbb{Z}_p$ has exactly p irreducible complex representations which are all one-dimensional and are generated by multiplication of a complex number by $\bar{\rho}^k$, $0 \leq k < p$ (Miller, 1972). When restricted to real spaces and $\rho^k \notin \mathbb{R}$, the multiplications by $\bar{\rho}^k$ and $\rho^k = \bar{\rho}^{p-k}$ can be combined to form a two-dimensional real irreducible representation space, in which the generator of $\mathbb{Z}_p$ acts by rotation of vectors by the angle $2\pi k/p$. For the representation of $\mathbb{Z}_p$ in the full space of p-dimensional row vectors generated by $\mathbf{P}$, the rows in V^* are (complex) basis vectors for these irreducible subspaces, and those basis vectors with eigenvalues $\bar{\rho}^k$ for which $\hat{\Sigma}$ has a nonzero column span the irreducible subspaces in span$\{\mathrm{R}(\Sigma)\}$. (Real bases in case of $\bar{\rho}^k \notin \mathbb{R}$ are obtained by taking real and imaginary parts of these vectors, but we prefer to use the complex basis vectors.) Likewise, the non-zero columns of $\hat{\Sigma}$ form complex bases of the irreducible subspaces of the $\mathbb{Z}_p$-representation generated by $\mathbf{J}$ in span$\{\mathrm{C}(\Sigma)\}$. We note that, given a row-vector $x \in \mathbb{C}^p$, xV is the discrete Fourier transform of x, and the components of xV are the expansion coefficients of x represented by the basis vectors in V^*/p.

Closely related to the group $\mathbb{Z}_p$ are the p-th roots of unity, which in turn are intimately related to the cyclotomic polynomials. In Sections 3-5 we will make some use of these polynomials and therefore summarize their basic properties in the appendix.

Given a row-vector $x \in \mathbb{R}^p$, the orbit of x under $\mathbf{P}$ is defined as the the set $\{x, x\mathbf{P}, \ldots, x\mathbf{P}^{p-1}\}$. For generic $x \in \mathbb{R}^p$, this set is a basis of $\mathbb{R}^p$, however the set of binary row vectors, $x = \eta$, is finite and the orbit of η, which we call a loop, may span only a proper subspace of $\mathbb{R}^p$. In the next section we classify (admissible) cycles according to the decomposition of $\mathrm{R}(\Sigma)$ into sets of rows belonging to different loops. To pursue this, we will introduce a concept of irreducibility that differs from the standard group-theoretical version above.

4. Classification of Cycles

4.1. Simple Cycles

Definition 2. Let $\eta = (\eta^1, \eta^2, \ldots, \eta^p)$ be a p-dimensional binary row vector. The set

$$\{\eta_\nu : \eta_\nu = \eta \mathbf{P}^\nu, \nu = 0, 1, 2, \ldots, p-1\},$$

is called a *loop* and is denoted by $\mathcal{L}_\eta$.

Remark 2. For any loop $\mathcal{L}_\eta$, $|\mathcal{L}_\eta| \le p$. More precisely, $|\mathcal{L}_\eta| = m$, where m is a factor of p. In particular, if $\eta = (+, +, \ldots, +)$, then $|\mathcal{L}_\eta| = 1$, as $\eta \mathbf{P} = \eta$.

Definition 3. A cycle Σ is called *simple*, if its row vectors are from one loop generated by some row vector η, i.e.,

$$\mathrm{R}(\Sigma) \subseteq \mathcal{L}_\eta. \tag{17}$$

A cycle Σ is *composite*, if it is not simple.

Definition 4. Let Σ be a cycle. The set $G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\}$ is said to be a *generator* of Σ, if

$$\mathcal{L}_{\eta_i} \cap \mathcal{L}_{\eta_j} = \emptyset, \ \forall i \neq j, \tag{18}$$

and

$$G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\} \subseteq \mathrm{R}(\Sigma) \subseteq \bigcup_{i=1}^{q} \mathcal{L}_{\eta_i}. \tag{19}$$

Note that any vector in $\mathcal{L}_{\eta_i} \cap \mathrm{R}(\Sigma)$ can be chosen as generator instead of η_i in G_Σ, that is, the generators are unique up to cyclic permutations and the condition to be vectors in $\mathrm{R}(\Sigma)$. In particular, for simple cycles there is only one generator, $|G_\Sigma| = 1$, and every row in $\mathrm{R}(\Sigma)$ can be chosen for this generator. A simple criterion for admissibility is the following.

Proposition 2. *A cycle Σ is admissible if $\mathcal{L}_\eta \subset \operatorname{span}\{\mathrm{R}(\Sigma) \cap \mathcal{L}_\eta\}$ for every $\eta \in G_\Sigma$.*

Proof. This follows immediately from the fact that, under the given hypothesis, every row in $\mathrm{R}(\Sigma\mathbf{P})$ can be represented as linear combination of a subset of rows in $\mathrm{R}(\Sigma)$. □

Definition 5. Let η be any row vector. The *rank* of η is defined as the dimension of the vector space spanned by the row vectors in the loop generated by η, i.e.,

$$\operatorname{rank}(\eta) = \dim \operatorname{span}\{\mathcal{L}_\eta\}. \tag{20}$$

Theorem 2. *Let Σ be a simple cycle generated by η, i.e., $\eta \in \mathrm{R}(\Sigma) \subseteq \mathcal{L}_\eta$. Then Σ is admissible, if and only if*

$$\operatorname{rank}(\Sigma) = \operatorname{rank}(\eta). \tag{21}$$

Proof. Suppose $\operatorname{rank}(\Sigma) = \operatorname{rank}(\eta)$. Then $\operatorname{span}\{\mathrm{R}(\Sigma)\} = \operatorname{span}\{\mathcal{L}_\eta\}$ as $\mathrm{R}(\Sigma) \subseteq \mathcal{L}_\eta$. Since $\mathbf{P}$ is nonsingular and $\mathcal{L}_\eta\mathbf{P} = \mathcal{L}_\eta$, it follows that $\operatorname{span}\{\mathrm{R}(\Sigma\mathbf{P})\} = \operatorname{span}\{\mathcal{L}_\eta\}$, hence Σ is admissible. Conversely, suppose that Σ is admissible, i.e. $\operatorname{span}\{\mathrm{R}(\Sigma)\} = \operatorname{span}\{\mathrm{R}(\Sigma\mathbf{P}^m)\}$ for all $m \in \mathbb{N}$. Assume $\operatorname{rank}(\Sigma) < \operatorname{rank}(\eta)$. Then there exists $\hat{\eta} \in \mathcal{L}_\eta$ with $\hat{\eta} \notin \operatorname{span}\{\mathrm{R}(\Sigma)\}$. Let $\hat{\eta} = \eta\mathbf{P}^\mu$ for some $0 < \mu < p$. Since $\eta \in \mathrm{R}(\Sigma)$, it follows that

$$\hat{\eta} \in \mathrm{R}(\Sigma\mathbf{P}^\mu) \subset \operatorname{span}\{\mathrm{R}(\Sigma\mathbf{P}^\mu)\} = \operatorname{span}\{\mathrm{R}(\Sigma)\},$$

which contradicts $\hat{\eta} \notin \operatorname{span}\{\mathrm{R}(\Sigma)\}$. □

Remark 3. In general, although $\mathbf{P}$ preserves the rank of any cycle Σ, the vector space spanned by the row vectors of Σ may not be invariant under $\mathbf{P}$. For simple cycles, the condition (21) guarantees that the vector space spanned by the rows of Σ is invariant under $\mathbf{P}$, and hence guarantees the admissibility of Σ. The condition (21) will be referred to as *admissibility condition* for simple cycles.

4.2. Separable Composite Cycles

In order to generalize the class of simple admissible cycles to the class of separable composite cycles, we first introduce the concept of decomposability of the row space of a cycle into irreducible subspaces.

Definition 6. Let Σ be an admissible cycle of period p, and let $\mathcal{U} = \operatorname{span}\{\mathrm{R}(\Sigma)\}$. Note that admissibility implies $\mathcal{U}\mathbf{P} = \mathcal{U}$, i.e. $\mathcal{U}$ is invariant under $\mathbf{P}$. Let $\mathcal{V} \subseteq \mathcal{U}$ be a subspace of $\mathcal{U}$ and assume $\mathcal{V}\mathbf{P} = \mathcal{V}$ and $\mathrm{R}(\Sigma) \cap \mathcal{V} \neq \emptyset$.

(a) The subspace $\mathcal{V}$ is called *reducible* if there exists a proper subspace $\mathcal{W}$ of $\mathcal{V}$ such that $\mathcal{W}\mathbf{P} = \mathcal{W}$ and $\mathrm{R}(\Sigma) \cap \mathcal{W} \neq \emptyset$.
(b) The subspace $\mathcal{V}$ is said to be *decomposable*, if $\mathcal{V}$ has the direct sum decomposition

$$\mathcal{V} = \mathcal{V}_1 \oplus \mathcal{V}_2 \oplus \cdots \oplus \mathcal{V}_n, \tag{22}$$

where $n \geq 2$, $\mathcal{V}_i$ is invariant under $\mathbf{P}$ and $\mathcal{V}_i \cap \mathrm{R}(\Sigma) \neq \emptyset$ for every $1 \leq i \leq n$. The subspace $\mathcal{V}$ is said to be *indecomposable*, if it is not decomposable. If $\mathcal{V}_i$ in (22) is indecomposable for every i, then (22) is called a *complete decomposition* of $\mathcal{V}$.
(c) The vector space $\mathcal{U}$ is called *semisimple*, if $\mathcal{U}$ is the direct sum of irreducible subspaces in the sense of (a). Note that semisimplicity of $\mathcal{U}$ includes the case where $\mathcal{U}$ is irreducible, in which case we call $\mathcal{U}$ simple.

Remark 4. We emphasize that, because our purpose is to study the structure of the invariant subspaces spanned by the row vectors of Σ, the concepts of *reducibility* and *decomposability* introduced in Definition 6 are slightly different from the standard definitions used in the representation theory of finite groups (we require that each subspace contains a row vector of Σ). An irreducible/indecomposable invariant subspace in the sense of Definition 6 may be reducible/decomposable in terms of the standard definitions of representation theory applied to the cyclic group $\mathbb{Z}_p$ generated by $\mathbf{P}$.

It is clear that if Σ is simple and admissible, then $\mathcal{U}$ is simple and consequently indecomposable, as $\eta \in \mathcal{U}_i$ implies that $\eta\mathbf{P}^k \in \mathcal{U}_i$ for every $k \in \mathbb{N}$. However, the converse is not necessarily true. In the next example, we show that the vector space spanned by the row vectors of a composite cycle may be reducible but not decomposable.

Example 1. Consider

$$\Sigma = \begin{pmatrix} + & + & + & - & - & - \\ + & + & - & - & - & + \\ + & - & - & - & + & + \\ + & - & + & - & + & - \end{pmatrix},$$

and let $\eta_j = \mathrm{row}_j(\Sigma)$. Clearly, Σ is a composite cycle, as it is generated by $\{\eta_1, \eta_4\}$. Let $\mathcal{U} = \mathrm{span}\{\mathrm{R}(\Sigma)\}$, $\mathcal{U}_1 = \mathrm{span}\{\mathcal{L}_{\eta_1}\}$ and $\mathcal{U}_2 = \mathrm{span}\{\mathcal{L}_{\eta_4}\}$. Since $\eta_4 = \eta_1 - \eta_2 + \eta_3$, i.e. $\eta_4 \in \mathcal{U}_1$, we have that $\mathcal{U}_2 \subset \mathcal{U}_1 = \mathcal{U}$. Moreover, $\mathrm{R}(\Sigma\mathbf{P}) = \{\eta_2, \eta_3, -\eta_1, -\eta_4\}$ implies $\mathcal{U}\mathbf{P} = \mathcal{U}$, hence Σ is admissible. Since both $\mathcal{U}_1$ and $\mathcal{U}_2$ are invariant under $\mathbf{P}$ it follows that $\mathcal{U}$ is reducible, however, $\mathcal{U}$ is not decomposable.

Proposition 3. *Let* Σ *be an admissible cycle with generator* $G_\Sigma = \{\eta_1, \ldots, \eta_q\}$*. Assume* $\mathcal{U} = \mathrm{span}\{\mathrm{R}(\Sigma)\}$ *is semisimple and let* $\mathcal{U} = \mathcal{U}_1 \oplus \mathcal{U}_2 \oplus \cdots \oplus \mathcal{U}_n$ *be a decomposition of* $\mathcal{U}$ *into irreducible subspaces. Then* $n \leq q$ *and there exists a subset* $\{i_1, \ldots, i_n\} \subseteq \{1, \ldots, q\}$ *such that* $\mathcal{U}_j = \mathrm{span}\{\mathcal{L}_{\eta_{i_j}}\}$ *for* $1 \leq j \leq n$*. Moreover, if* $\mathrm{span}\{\mathcal{L}_{\eta_i}\} \neq \mathrm{span}\{\mathcal{L}_{\eta_j}\}$ *for every* $i, j \in \{1, \ldots, q\}$ *with* $i \neq j$*, then* $n = q$*.*

Proof. Let $\eta \in G_\Sigma$ and let $\mathcal{V} \in \{\mathcal{U}_j | 1 \leq j \leq n\}$ be the subspace in the decomposition of $\mathcal{U}$ that contains η. Invariance of $\mathcal{V}$ implies $\eta\mathbf{P} \in \mathcal{V}$, hence $\eta\mathbf{P}^2 \in \mathcal{V}$ and by induction $\mathcal{L}_\eta \subset \mathcal{V}$, thus $\mathrm{span}\{\mathcal{L}_\eta\} \subset \mathcal{V}$. Since $\mathrm{span}\{\mathcal{L}_\eta\}$ is invariant, $\mathrm{span}\{\mathcal{L}_\eta\} \cap \mathrm{R}(\Sigma) \neq \emptyset$ and $\mathcal{V}$ is irreducible, it follows that $\mathcal{V} = \mathrm{span}\{\mathcal{L}_\eta\}$, and there exists no $\eta' \in G_\Sigma$, $\eta' \neq \eta$, such that $\mathrm{span}\{\mathcal{L}_{\eta'}\}$ is a proper subspace of $\mathrm{span}\{\mathcal{L}_\eta\}$ and vice versa. Thus, for $i \neq j$, either $\mathrm{span}\{\mathcal{L}_{\eta_i}\} \cap \mathrm{span}\{\mathcal{L}_{\eta_j}\} = \{\mathbf{0}\}$ or $\mathrm{span}\{\mathcal{L}_{\eta_i}\} = \mathrm{span}\{\mathcal{L}_{\eta_j}\}$. It follows that there exists $i_1, \ldots, i_n \in \{1, \ldots, q\}$, $i_j \neq i_k$ if $j \neq k$, such that

$$\mathrm{span}\{\mathrm{R}(\Sigma)\} = \mathrm{span}\{\bigcup_{i=1}^{q} \mathcal{L}_{\eta_i}\} = \bigoplus_{j=1}^{n} \mathrm{span}\{\mathcal{L}_{\eta_{i_j}}\}.$$

□

Example 2. Let $\eta_1 = (+, +, -, +, +, -)$, $\eta_2 = -\eta_1$ and $\eta_3 = (+, +, +, -, -, -)$. Let Σ be the 9×6-cycle defined by

$$\Sigma = (\Sigma_1^T, \Sigma_2^T, \Sigma_3^T)^T,$$

where $\Sigma_j = (\eta_j^T, (\eta_j\mathbf{P})^T, (\eta_j\mathbf{P}^2)^T)^T$ for $j = 1, 2, 3$. Then $G_\Sigma = \{\eta_1, \eta_2, \eta_3\}$, $\mathrm{span}\{\mathrm{R}(\Sigma)\} = \mathrm{span}\{\mathcal{L}_{\eta_1}\} \oplus \mathrm{span}\{\mathcal{L}_{\eta_3}\}$, and $\mathrm{span}\{\mathcal{L}_{\eta_2}\} = \mathrm{span}\{\mathcal{L}_{\eta_1}\}$. Clearly, $\mathrm{span}\{\mathcal{L}_{\eta_1}\}$ and $\mathrm{span}\{\mathcal{L}_{\eta_3}\}$ are irreducible, thus $\mathrm{span}\{\mathrm{R}(\Sigma)\}$ is semisimple. Likewise, for the cycle $\Sigma = (\Sigma_1^T, \Sigma_2^T)$, $\mathrm{span}\{\mathrm{R}(\Sigma)\} = \mathrm{span}\{\mathcal{L}_{\eta_i}\}$, $i = 1, 2$, hence $\mathrm{span}\{\mathrm{R}(\Sigma)\}$ is simple.

In general, the vector space $\mathcal{U} = \mathrm{span}\{\mathrm{R}(\Sigma)\}$ of an arbitrary composite cycle Σ may have subspaces which are not invariant or do not contain any binary row vector of Σ, or both. By contrast, if $\mathcal{U}$ is semisimple, $\mathcal{U}$ can be decomposed into irreducible subspaces corresponding to the loops of their generators, but some of these subspaces may coincide. This coincidence is still considered as a degeneracy (see Section 4.3), which we exclude in the class of separable cycles introduced next.

Definition 7. Let Σ be a composite cycle with generator G_Σ, i.e. $|G_\Sigma| \geq 2$. We call Σ *separable*, if $\operatorname{span}\{\mathrm{R}(\Sigma)\}$ is semisimple and $\operatorname{span}\{\mathcal{L}_\eta\} \neq \operatorname{span}\{\mathcal{L}_{\eta'}\}$ for any $\eta, \eta' \in G_\eta$ with $\eta \neq \eta'$. If Σ is not separable, Σ is said to be *inseparable*.

Note that the hypotheses for a cycle Σ to be separable require that $\mathcal{U} = \operatorname{span}\{\mathrm{R}(\Sigma)\}$ is invariant under $\mathbf{P}$, i.e. separable cycles are a priori admissible.

Theorem 3. ***(Separability Condition for Composite Cycles)*** *Let Σ be a composite cycle with generator $G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\}$. Then Σ is separable, if and only if*

$$\operatorname{rank}(\Sigma) = \sum_{i=1}^{q} \operatorname{rank}(\eta_i). \tag{23}$$

Proof. If Σ is separable, (23) follows directly from Proposition 3 and Definition 7.

Conversely, suppose (23) holds. Since

$$\operatorname{span}\{\mathrm{R}(\Sigma)\} = \operatorname{span}\{(\bigcup_{i=1}^{q} \mathcal{L}_{\eta_i}) \cap \mathrm{R}(\Sigma)\} \subseteq \operatorname{span}\{\bigcup_{i=1}^{q} \mathcal{L}_{\eta_i}\},$$

(23) implies that $\operatorname{span}\{\mathcal{L}_{\eta_i} \cap \mathrm{R}(\Sigma)\} = \operatorname{span}\{\mathcal{L}_{\eta_i}\}$ for each $1 \leq i \leq q$, $\operatorname{span}\{\mathcal{L}_{\eta_i}\} \cap \operatorname{span}\{\mathcal{L}_{\eta_j}\} = \{\mathbf{0}\}$, if $i \neq j$, and hence

$$\operatorname{span}\{\mathrm{R}(\Sigma)\} = \bigoplus_{i=1}^{q} \operatorname{span}\{\mathcal{L}_{\eta_i}\}.$$

It follows that each $\operatorname{span}\{\mathcal{L}_{\eta_i}\}$ is irreducible, thus Σ is separable according to Definitions 6 and 7. $\square$

Example 3. Consider

$$\Sigma = \begin{pmatrix} + & + & + & + & - & - & - & - \\ + & + & + & - & - & - & - & + \\ + & + & - & - & - & - & + & + \\ + & - & - & - & - & + & + & + \\ + & + & - & - & + & + & - & - \\ + & - & - & + & + & - & - & + \\ + & - & + & - & + & - & + & - \end{pmatrix}.$$

We have that $G_\Sigma = \{\eta_1, \eta_5, \eta_7\}$, where $\eta_j = \mathrm{row}_j(\Sigma)$, $1 \leq j \leq 7$. It is easy to see that $\mathcal{U}_1 = \mathrm{span}\{\mathcal{L}_{\eta_1}\}$, $\mathcal{U}_2 = \mathrm{span}\{\mathcal{L}_{\eta_5}\}$, and $\mathcal{U}_3 = \mathrm{span}\{\mathcal{L}_{\eta_7}\}$ intersect trivially, hence

$$\mathcal{U} = \mathrm{span}(\mathrm{R}(\Sigma)) = \mathcal{U}_1 \oplus \mathcal{U}_2 \oplus \mathcal{U}_3,$$

which implies that Σ is separable, and hence admissible.

4.3. Inseparable Composite Cycles

By Definition 7, inseparability of a composite cycle Σ happens in two different cases. In the first case, the vector space $\mathcal{U} = \mathrm{span}\{\mathrm{R}(\Sigma)\}$ has a reducible but indecomposable invariant subspace, which entirely contains another invariant subspace as a subspace (see Example 1). This includes the case where $\mathcal{U}$ is semisimple and $\mathrm{span}\{\mathcal{L}_\eta\} = \mathrm{span}\{\mathcal{L}_{\eta'}\}$ for two different generators η and η' (see Example 2). In the second case, the vector space $\mathcal{U}$ has two or more indecomposable (reducible or not) invariant subspaces sharing a nontrivial intersection as common proper subspaces. We now discuss the admissibility of these two types of inseparable composite cycles.

Definition 8. Let $G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\}$ be a generator of a cycle Σ. A subset $EG_\Sigma = \{\epsilon_1, \epsilon_2, \ldots, \epsilon_r\}$, $r \leq q$, of G_Σ is called an *essential generator* of Σ, if EG_Σ is minimal in the sense that

(a) $\mathrm{span}\{\mathrm{R}(\Sigma)\} \subseteq \mathrm{span}\{\bigcup_{i=1}^{r} \mathcal{L}_{\epsilon_i}\}$;

(b) for any ϵ_i, $\epsilon_j \in EG_\Sigma$ with $i \neq j$, $\mathrm{span}(\mathcal{L}_{\epsilon_i}) \cap \mathrm{span}(\mathcal{L}_{\epsilon_j})$ is a proper subspace of both $\mathrm{span}(\mathcal{L}_{\epsilon_i})$ and $\mathrm{span}(\mathcal{L}_{\epsilon_j})$;

(c) for every $\eta_i \in G_\Sigma$, if $\eta_i \notin \mathrm{span}\{\bigcup_{j=1, j\neq i}^{q} \mathcal{L}_{\eta_j}\}$, then $\eta_i \in EG_\Sigma$.

Note that an admissible cycle may have different sets of essential generators, i.e. EG_Σ is in general not unique. Proposition 2 can be directly rephrased in terms of essential generators.

Proposition 4. *A cycle Σ is admissible, if for any essential generator EG_Σ, $\mathcal{L}_\epsilon \subset \mathrm{span}\{\mathrm{R}(\Sigma) \cap \mathcal{L}_\epsilon\}$ for every $\epsilon \in EG_\Sigma$. Conversely, if Σ is admissible and $EG_\Sigma = \{\epsilon_1, \ldots, \epsilon_r\}$ is an essential generator of Σ, then*

$$\mathrm{rank}(\Sigma) \leq \sum_{i=1}^{r} \mathrm{rank}(\epsilon_i) \leq |\mathrm{R}(\Sigma)| \leq N. \tag{24}$$

Remark 5. The condition (b) in Definition 8 includes three cases.
(a) For every $i \neq j$, $\mathrm{span}\{\mathcal{L}_{\epsilon_i}\} \cap \mathrm{span}\{\mathcal{L}_{\epsilon_j}\} = \{\mathbf{0}\}$, and $q = r$. Composite cycles in this case are separable.
(b) For every $i \neq j$, $\mathrm{span}\{\mathcal{L}_{\epsilon_i}\} \cap \mathrm{span}\{\mathcal{L}_{\epsilon_j}\} = \{\mathbf{0}\}$, but $q < r$. Composite cycles in this case are inseparable and degenerate. In this case $\mathcal{U}$ may be semisimple or not, and has the complete decomposition $\mathcal{U} = \bigoplus_{i=1}^{q} \mathrm{span}\{\mathcal{L}_{\epsilon_i}\}$ (see Proposition 3). Accordingly, we have that a degenerately inseparable composite cycle Σ is admissible, if and only if $\mathrm{rank}(\Sigma) = \sum_{i=1}^{q} \mathrm{rank}(\epsilon_i)$. This generalizes separable cycles and includes, for example, the case where for some $\eta \in \mathrm{R}(\Sigma)$ also $-\eta \in \mathrm{R}(\Sigma)$, but $\mathcal{L}_\eta \cap \mathcal{L}_{-\eta} = \emptyset$ (see Example 2).
(c) For some $i \neq j$, $\mathrm{span}\{\mathcal{L}_{\epsilon_i}\} \cap \mathrm{span}\{\mathcal{L}_{\epsilon_j}\}$ is a nontrivial proper subspace of both invariant subspaces $\mathrm{span}\{\mathcal{L}_{\epsilon_i}\}$ and $\mathrm{span}\{\mathcal{L}_{\epsilon_j}\}$. Composite cycles in this case are genuinely inseparable. This type of cycles is more complicated than the other two. We next study the structure of this type of cycles, and establish an admissibility condition.

Proposition 5. *Let η and $\tilde{\eta}$ be two p-dimensional row vectors. If $\eta \in \mathrm{span}\{\mathcal{L}_{\tilde{\eta}}\}$, then $\mathrm{span}\{\mathcal{L}_\eta\} \subseteq \mathrm{span}\{\mathcal{L}_{\tilde{\eta}}\}$.*

Proof. If $\eta \in \mathcal{L}_{\tilde{\eta}}$, we are done. Suppose $\eta \in \mathrm{span}\{\mathcal{L}_{\tilde{\eta}}\}$, i.e., $\eta = \sum_{\nu=1}^{p} \alpha_\nu \tilde{\eta}\mathbf{P}^\nu$, $\alpha_\nu \in \mathbb{R}$, but $\eta \notin \mathcal{L}_{\tilde{\eta}}$, then for every $t \in \{1, 2, \ldots, p\}$, $\eta\mathbf{P}^t = \left(\sum_{\nu=1}^{p} \alpha_\nu \tilde{\eta}\mathbf{P}^\nu\right)\mathbf{P}^t = \sum_{\nu=1}^{p} (\alpha_\nu \tilde{\eta}\mathbf{P}^{\nu+t})$, i.e., $\mathcal{L}_\eta \subseteq \mathrm{span}\{\mathcal{L}_{\tilde{\eta}}\}$, hence $\mathrm{span}\{\mathcal{L}_\eta\} \subseteq \mathrm{span}\{\mathcal{L}_{\tilde{\eta}}\}$. □

Remark 6. Proposition 5 tells that if a row vector is in the vector space spanned by the loop generated by another row vector of the same dimension, then the vector space spanned by the loop generated by this row vector is a subspace of the vector space spanned by the other one. Since for any genuinely inseparable composite cycle, at least two indecomposable invariant subspaces intersect nontrivially, it is natural to ask:
(a) Does there exist a row vector such that this nontrivial intersection is spanned by the loop generated by it?
(b) If this row vector exists, can it be $\{-1, 1\}$-valued?

As we will see below in Proposition 6, the answer to the first question is affirmative, however, it remains unclear whether there always exists a binary row vector such that the loop generated by it spans the nontrivial intersection of two indecomposable invariant subspaces. The approach we will use in the proof of Proposition 6 only guarantees the existence of a genuine row vector, which may or may not be binary.

Let V be defined as in Theorem 1, i.e. $V = (v^{(0)}, v^{(1)}, \ldots, v^{(p-1)})$, where $v^{(k)} = (1, \rho^k, \rho^{2k}, \ldots, \rho^{(p-1)k})^T$ and $\rho = e^{2\pi \mathbf{i}/p}$.

Definition 9. A row vector η is said to *annihilate* the column $v^{(k)}$ of V, if $\eta v^{(k)} = 0$, i.e. the two vectors are orthogonal.

Note that, since $v^{(k)}$ is an eigenvector of $\mathbf{P}$ and all eigenvalues of $\mathbf{P}$ are nonzero, η annihilates $v^{(k)}$ if and only if $\eta \mathbf{P}^\nu$ annihilates $v^{(k)}$ for every $\nu \in \mathbb{Z}$. We need the following fact about the eigenvectors and eigenvalues of circulant matrices, see, e.g., Lancaster & Tismenetsky (1985).

Lemma 1. *Let $\eta = (\eta^1, \ldots, \eta^p)$ be an arbitrary real and nonzero row vector, and let Σ_η be the $p \times p$-matrix defined by $\mathrm{row}_j(\Sigma_\eta) = \eta(\mathbf{P}^T)^{(j-1)}$ for $1 \leq j \leq p$. Then $V^*\Sigma_\eta = \Lambda_\eta V^*$, where $\Lambda_\eta = \mathrm{diag}(\lambda_{\eta,1}, \ldots, \lambda_{\eta,p})$ with $\lambda_{\eta,k} = \sum_{j=1}^{p} \eta^j \rho^{(j-1)(k-1)}$.*

Extending Definition 2 to non-binary real row vectors and noting that $\mathrm{R}(\Sigma_\eta) = \mathcal{L}_\eta$, an immediate consequence of Lemma 1 is the following:

Corollary 1. *Assume that $\eta v^{(j)} \neq 0$ if and only if $j \in \{k_1, \ldots, k_s\} \subset \mathbb{Z}_p$, where $\mathbb{Z}_p = \{0, 1, 2, \ldots, p-1\}$. Then $\{v^{(k_1)*}, \ldots, v^{(k_s)*}\}$ is a (complex) basis for $\mathrm{span}\{\mathcal{L}_\eta\}$.*

Proposition 6. *Let Σ be a cycle with essential generator $EG_\Sigma = \{\epsilon_1, \ldots, \epsilon_r\}$. Assume that for some $i \neq j$ the indecomposable subspaces $\mathcal{U}_i = \mathrm{span}(\mathcal{L}_{\epsilon_i})$ and $\mathcal{U}_j = \mathrm{span}(\mathcal{L}_{\epsilon_j})$ intersect nontrivially, and $\mathcal{U}_i \cap \mathcal{U}_j$ is a proper subspace of both $\mathcal{U}_i$ and $\mathcal{U}_j$. Then there exists a row vector η such that $\mathcal{U}_i \cap \mathcal{U}_j = \mathrm{span}\{\mathcal{L}_\eta\}$.*

Proof. Assume that $\epsilon_i v^{(k)} \neq 0$ and $\epsilon_j v^{(k)} \neq 0$ if and only if $k \in K_i \subset \mathbb{Z}_p$ and $k \in K_j \subset \mathbb{Z}_p$, respectively. Assume further that $K \equiv K_i \cap K_j = \{k_1, \ldots, k_s\}$. According to Corollary 1, $\{v^{(k_1)*}, \ldots, v^{(k_s)*}\}$ is a basis for $\mathcal{U}_i \cap \mathcal{U}_j$. Let $p_i(x)$ and $p_j(x)$, $x \in \mathbb{C}$, be the polynomials $p_i(x) = \epsilon_i \mathbf{x}$, $p_j(x) = \epsilon_j \mathbf{x}$, where $\mathbf{x} = (1, x, x^2, \ldots, x^{p-1})^T$. Since the row vectors defined by the coefficients of $p_i(x)$ and $p_j(x)$ annihilate exactly the $v^{(k)}$ with $k \in \mathbb{Z}_p \backslash K_i$ and $k \in \mathbb{Z}_p \backslash K_j$, respectively, $p_i(x)$ and $p_j(x)$ contain the minimal polynomials of ρ^k for every $k \in \mathbb{Z}_p \backslash K_i$ and $k \in \mathbb{Z}_p \backslash K_j$, respectively, as factors. Multiplying these factors yields a polynomial $p_{ij}(x)$ of degree $\leq p-1$ with $p_{ij}(\rho^k) = 0$ for every

$k \in \mathbb{Z}_p \backslash K$ and $p_{ij}(\rho^k) \neq 0$ for every $k \in K$. Set $p_0(x) = p_{ij}(x)$ if the degree of $p_{ij}(x)$ is $p-1$. If the degree of $p_{ij}(x)$ is $< p-1$, set $p_0(x) = p_{ij}(x)\tilde{p}_{ij}(x)$, where $\tilde{p}_{ij}(x)$ is any polynomial such that $\tilde{p}_{ij}(\rho^k) \neq 0$ for every $k \in K$ and the degree of $p_0(x)$ is $p-1$. Let η be the row vector of coefficients of $p_0(x)$. Then η and $\eta\mathbf{P}^\nu$ for any $\nu \in \mathbb{Z}$ annihilate every $v^{(k)}$ for $k \in \mathbb{Z}_p\backslash K$, and $\eta v^{(k)} \neq 0$ for every $k \in K$, hence $\text{span}\{\mathcal{L}_\eta\} = \text{span}\{v^{(k_1)*}, \ldots, v^{(k_s)*}\} = \mathcal{U}_i \cap \mathcal{U}_j$. □

Example 4. In this example, we demonstrate how to find a row vector as claimed in Proposition 6 with the method described in the proof. Consider the composite cycle with $N = 10$ and $p = 18$ defined by $\Sigma = (\Sigma_1^T, \Sigma_2^T)^T$, where

$$\Sigma_1 = (\epsilon_1^T, (\epsilon_1\mathbf{P})^T, \ldots, (\epsilon_1\mathbf{P}^6)^T)^T, \Sigma_2 = (\epsilon_2^T, (\epsilon_2\mathbf{P})^T, (\epsilon_1\mathbf{P}^2)^T)^T,$$

and

$$\begin{aligned}\epsilon_1 &= (\ + \ + \ + \ + \ + \ + \ + \ - \ + \ - \ - \ - \ - \ - \ - \ - \ + \ - \ ),\\ \epsilon_2 &= (\ + \ + \ + \ - \ - \ - \ + \ + \ + \ - \ - \ - \ + \ + \ + \ - \ - \ - \ ).\end{aligned}$$

This is a genuinely inseparable composite cycle with $G_\Sigma = EG_\Sigma = \{\epsilon_1, \epsilon_2\}$. The polynomials $p_1(x)$ and $p_2(x)$ can be factorized as follows,

$$\begin{aligned}p_1(x) &= (1-x)(1+x+x^2)(1-x+x^2)(1+x^3+x^6)(1+2x+2x^2+x^3+x^6),\\ p_2(x) &= (1-x)(1+x+x^2)(1-x^3+x^6)(1+x^3+x^6)(1+x-x^2).\end{aligned}$$

The factors $1-x$, $1+x+x^2$, $1+x^3+x^6$, $1-x+x^2$ and $1-x^3+x^6$ are cyclotomic factors (see e.g. Dummit & Foote (2004)), and the sum of their degrees happens to be 17. Multiplying them out gives

$$p_0(x) = \sum_{j=0}^{17} (-1)^j x^j,$$

thus the row vector η constructed in the proof of Proposition 6 is obtained as the binary vector with alternating signs, $\eta = (+, -, +, -, \ldots, +, -)$, and $\text{span}\{\mathcal{L}_\eta\} = \text{span}\{\eta\}$. One can easily verify that $\text{span}\{\eta\} = \text{span}\{\mathcal{L}_{\epsilon_1}\} \cap \text{span}\{\mathcal{L}_{\epsilon_2}\}$.

Based on their structural features and using a simple inclusion-exclusion argument, an admissibility condition for inseparable composite cycles can be formulated as follows.

Theorem 4. ***(Admissibility Condition for Inseparable Composite Cycles)*** *Let Σ be a cycle with essential generator $EG_\Sigma = \{\epsilon_1, \ldots, \epsilon_r\}$. Then Σ is admissible if and only if*

$$\begin{aligned} \operatorname{rank}(\Sigma) &= \sum_{i=1}^{r} \operatorname{rank}(\epsilon_i) - \sum_{i,j;i\neq j}^{r} \dim(\operatorname{span}\{\mathcal{L}_{\epsilon_i}\} \cap \operatorname{span}\{\mathcal{L}_{\epsilon_j}\}) \\ &+ \sum_{i,j,k;i\neq j\neq k}^{r} \dim(\operatorname{span}\{\mathcal{L}_{\epsilon_i}\} \cap \operatorname{span}\{\mathcal{L}_{\epsilon_j}\} \cap \operatorname{span}\{\mathcal{L}_{\epsilon_k}\}) \qquad (25) \\ &- \cdots \pm \dim(\bigcap_{i=1}^{r} \operatorname{span}\{\mathcal{L}_{\epsilon_i}\}). \end{aligned}$$

Remark 7. The admissibility condition (25) is valid for any cycle, and the conditions (23) and (21) for separability and admissibility of simple cycles, respectively, can be thought of as special cases thereof.

Example 5. Let Σ be the cycle from Example 4. One can easily verify that $\operatorname{rank}(\epsilon_1) = 7$ and $\operatorname{rank}(\epsilon_2) = 3$. Since $\dim(\operatorname{span}\{\mathcal{L}_{\epsilon_1}\} \cap \operatorname{span}\{\mathcal{L}_{\epsilon_2}\}) = 1$, this cycle is admissible.

5. Network Topology

To simplify the discussion, we exclude in this section multiple appearances of a binary row vector in a cycle, that is, we consider only cycles Σ with $|\mathrm{R}(\Sigma)| = N$.

The classification of cycles Σ in Section 4 was based on the decomposition of $\mathrm{R}(\Sigma)$ into subsets of rows associated with disjoint loops. It is, therefore, natural to identify the neurons corresponding to the same loop with a cluster. However, if a cycle has fewer essential generators than generators, the row vectors of a non-essential generator must be combined with one or more essential generators and, moreover, there may be several choices for essential generators. We therefore make the simplifying assumption that all generators are essential generators. For admissible cycles Σ this means that for any two distinct generators η_1, η_2, the intersection of their spaces $\operatorname{span}\{\mathcal{L}_{\eta_j} \cap \mathrm{R}(\Sigma)\}$, $j = 1, 2$, is a proper subspace of both of them ($\{\mathbf{0}\}$ if the cycle is separable). An immediate consequence of this assumption is that

$$\operatorname{rank}(\Sigma) \leq \sum_{j=1}^{q} \operatorname{rank}(\eta_j) \leq N, \qquad (26)$$

if Σ is admissible and $G_\Sigma = \{\eta_1, \ldots, \eta_q\}$. The clusters are isolated if and only if Σ is separable. If Σ is inseparable, some of the clusters are connected.

Regarding the connectivity within a cluster, linear dependences among its row vectors will prevent any special structure. We call cycles for which such dependences do not occur minimal.

Definition 10. An admissible cycle Σ with generator $G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\}$ is *minimal*, if $EG_\Sigma = G_\Sigma$ and for every $1 \leq i \leq q$,

$$|\mathcal{L}_{\eta_i} \cap \mathrm{R}(\Sigma)| = \mathrm{rank}(\eta_i). \tag{27}$$

Remark 8. If Σ is a minimal simple or separable composite cycle, then Σ is of full row rank. If Σ is a minimal inseparable cycle, then the row vectors in $\mathrm{R}(\Sigma) \cap \mathcal{L}_{\eta_i}$ form a basis of $\mathrm{span}\{\mathcal{L}_{\eta_i}\}$ for every $\eta_i \in G_\Sigma$. Thus for any minimal cycle Σ, (26) holds and the inequalities become equalities if and only if Σ is simple or separable. In this case, Σ has full row rank and $\Sigma^+ = \Sigma^T(\Sigma\Sigma^T)^{-1}$, which implies that $\mathbf{J^0} = \Sigma\Sigma^+ = I$, the $N \times N$ identity matrix.

For any two $(N \times p)$-cycles Σ and Σ' with $\mathrm{R}(\Sigma) = \mathrm{R}(\Sigma')$, the cycle matrices are related to each other by $\Sigma' = \mathbf{Q}\Sigma$, where $\mathbf{Q}$ is an $N \times N$ permutation matrix. If in addition Σ is admissible with connectivity matrix $\tilde{\mathbf{J}}$, then Σ' is also admissible and has connectivity matrix $\tilde{\mathbf{J}}' = \mathbf{Q}\tilde{\mathbf{J}}\mathbf{Q}^{-1}$. Accordingly, if $\mathbf{u}$ is the state of the network with connectivity matrix $\tilde{\mathbf{J}}$, then $\mathbf{u}' = \mathbf{Qu}$ is the network state corresponding to $\tilde{\mathbf{J}}'$, and solutions $\mathbf{u}(t)$ and $\mathbf{u}'(t)$ of the corresponding differential equations are just permutations of each other as $\tanh(\mathbf{u}') = \tanh(\mathbf{Q}^{-1}\mathbf{u}) = \mathbf{Q}^{-1}\tanh(\mathbf{u})$.

Without loss of generality, we therefore may assume that a minimal cycle with generators $\eta_1, \ldots, \eta_q$ has the form

$$\Sigma = (\Sigma_1^T, \Sigma_2^T, \ldots, \Sigma_q^T)^T, \tag{28}$$

where $\mathrm{R}(\Sigma_j) \subseteq \mathcal{L}_{\eta_j}$, $1 \leq j \leq q$, and the vectors in Σ_j are sorted from top to bottom as $\eta_j, \eta_{j_1}, \eta_{j_2}, \ldots$, with $\eta_{j_i} = s_{j_i}\eta_j\mathbf{P}^{\nu_{j_i}}$, $s_{j_i} = 1$ if $-\eta_j \notin \mathcal{L}_{\eta_j}$ and $s_{j_i} \in \{-1, 1\}$ if $-\eta_j \in \mathcal{L}_{\eta_j}$, and $0 < \nu_{j_i} < \nu_{j_k}$ if $i < k$. We call this form the *standard form* of a minimal cycle.

The minimality requirement does not suffice in general to induce a special network topology within the clusters. We have to require in addition that the powers in the Σ_j are consecutive.

Definition 11. A minimal cycle Σ in standard form is said to be a *minimal consecutive cycle*, or briefly MC-cycle, if the powers of $\mathbf{P}$ in Σ_j above are consecutive, that is, $\nu_{j_i} = i$ for all $1 \leq i < \mathrm{rank}(\eta_j)$.

In order that Definition 11 is consistent with the minimality requirement, the rows in Σ_j must be linearly independent. The next proposition shows that this is indeed the case, where for simplicity we consider only the case $s_{j_i} = 1$.

Proposition 7. *Let* $\eta \neq 0$ *be any* p*-dimensional row vector with* $\mathrm{rank}(\eta) = k$*. Then the vectors* $\{\eta, \eta\mathbf{P}, \eta\mathbf{P}^2, \ldots, \eta\mathbf{P}^{k-1}\}$ *are linearly independent.*

Proof. Let s be the smallest positive integer such that $\{\eta, \eta\mathbf{P}, \ldots, \eta\mathbf{P}^{s-1}\}$ are linearly independent. Then $\eta\mathbf{P}^s$ is a linear combination of $\{\eta, \eta\mathbf{P}, \ldots, \eta\mathbf{P}^{s-1}\}$,

$$\eta\mathbf{P}^s = \sum_{\nu=0}^{s-1} \alpha_\nu \eta\mathbf{P}^\nu. \tag{29}$$

Right-multiplying this equation by $\mathbf{P}$ yields a representation of $\eta\mathbf{P}^{s+1}$ as linear combination of $\{\eta\mathbf{P}, \eta\mathbf{P}^2, \ldots, \eta\mathbf{P}^s\}$, and replacing $\eta\mathbf{P}^s$ in this representation by (29) shows that $\eta\mathbf{P}^{s+1}$ is also a linear combination of $\{\eta, \eta\mathbf{P}, \ldots, \eta\mathbf{P}^{s-1}\}$. By induction we find that, for any $0 \leq \nu < s - p$, $\eta\mathbf{P}^{s+\nu}$ is a linear combination of $\{\eta, \eta\mathbf{P}, \ldots, \eta\mathbf{P}^{s-1}\}$, hence this set is a basis for $\mathcal{L}_\eta$. $\square$

For simple MC-cycles there is only one cluster. In Subsection 5.1 we discuss the possible connectivity structures in such networks in some detail, including the possible values of N for a given p, and we also comment on the network topology of simple minimal but non-consecutive cycles. Semisimple MC-cycles consist of isolated clusters corresponding to the different loops in the cycle. Each of these loops forms a simple MC-cycle, and we just give an example in Subsection 5.2. Inseparable minimal (consecutive or non-consecutive) cycles are more complicated and will be discussed in Subsection 5.3. In Subsection 5.4 we demonstrate the effects of fewer essential generators than generators by two examples.

5.1. Simple MC-Cycles

5.1.1. Network Topology

According to Definition 11, a simple MC-cycle has the form

$$\Sigma = (\eta^T, s_1(\eta\mathbf{P})^T, s_2(\eta\mathbf{P}^2)^T, \ldots, s_{N-1}(\eta\mathbf{P}^{N-1})^T)^T, \tag{30}$$

with $\mathrm{rank}(\eta) = N \leq p$ and $s_i \in \{-1, 1\}$. Since the image of the last row vector of Σ under $\mathbf{P}$ is a linear combination of the row vectors of Σ, $\Sigma\mathbf{P}$ has the form $\Sigma\mathbf{P} = \mathbf{A}\Sigma$, where

$$\mathbf{A} = \begin{pmatrix} 0 & s_1 & 0 & \dots & 0 & 0 \\ 0 & 0 & s_1 s_2 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & s_{N-3}s_{N-2} & 0 \\ 0 & 0 & 0 & \dots & 0 & s_{N-2}s_{N-1} \\ a_1 & a_2 & a_3 & \dots & a_{N-1} & a_N \end{pmatrix}, \tag{31}$$

with $a_1, \dots, a_N \in \mathbb{R}$ and $a_1 \neq 0$. Moreover, since Σ has full row rank, $\Sigma^+ = \Sigma^T(\Sigma\Sigma^T)^{-1}$, which implies

$$\mathbf{J} = \Sigma\mathbf{P}\Sigma^+ = \mathbf{A}. \tag{32}$$

Equations (31) and (32) show that the network constructed from a simple MC-cycle consists of a feed-forward chain from the Nth neuron to the first neuron, and feedback to the Nth neuron from the subset of the neurons for which $a_i \neq 0$, which in any case includes the first neuron. If $a_j = 0$ for $j > 1$, then $a_1 = \pm 1$, and the network topology is that of a ring, with either excitatory ($a_1 = 1$, $\mathbf{J} = \mathbf{P}^T$ if all $s_j = 1$) or inhibitory connection ($a_1 = -1$) from neuron 1 to neuron N. Vectors of the form $\eta = (\sigma, \sigma)$ or $(\sigma, -\sigma)$ have $\mathrm{rank}(\eta) \leq p/2$, and if $\mathrm{rank}(\eta) = p/2 = N$ we have either of these two types of ring structures (see Subsection 5.1.3.).

Example 6. In Figure 2, A and B, we illustrate the topology of the networks constructed from the following two simple MC-cycles,

$$\Sigma = \begin{pmatrix} + & + & + & + & + & + & + & - & - & - & - & - & - & - \\ + & + & + & + & + & + & - & - & - & - & - & - & - & + \\ + & + & + & + & + & - & - & - & - & - & - & - & + & + \\ + & + & + & + & - & - & - & - & - & - & - & + & + & + \\ + & + & + & - & - & - & - & - & - & - & + & + & + & + \\ + & + & - & - & - & - & - & - & - & + & + & + & + & + \\ + & - & - & - & - & - & - & - & + & + & + & + & + & + \end{pmatrix},$$

and

$$\tilde{\Sigma} = \begin{pmatrix} + & + & - & + & - & - \\ + & - & + & - & - & + \\ - & + & - & - & + & + \\ + & - & - & + & + & - \\ - & - & + & + & - & + \end{pmatrix},$$

respectively. The cycle Σ has a "repeating block structure", $\Sigma = [B, -B]$, where B is the block consisting of the first 7 columns of Σ ($N = p/2 = 7$). This causes the image of the last row to be the negative copy of the first row, i.e. $\eta_7\mathbf{P} = -\eta_1$, where $\eta_i = \text{row}_i(\Sigma)$. It follows that $a_1 = -1$ and $a_i = 0$ for $i > 1$, thus the first neuron only sends an inhibitory feedback to the seventh neuron (Figure 2A). Similarly, for the cycle $\tilde{\Sigma}$, $N = p - 1 = 5$, and in this case the image of the last row is a linear combination of all other row vectors, $\tilde{\eta}_5\mathbf{P} = -\tilde{\eta}_1 - \tilde{\eta}_2 - \tilde{\eta}_3 - \tilde{\eta}_4 - \tilde{\eta}_5$, where $\tilde{\eta}_i = \text{row}_i(\tilde{\Sigma})$. Accordingly for this cycle $a_i = -1$ for every i, i.e. every neuron sends inhibitory feedback to the fifth neuron in the network (Figure 2B).

The two examples above demonstrate that the value of $N = \text{rank}(\eta)$ plays an important role for the network topology of simple MC-cycles. We discuss possible values of N for given cycle-lengths p in the next paragraph.

For minimal but non-consecutive cycles with $N < p - 1$ we can have "gaps" in the standard form which lead to feedforward chains interrupted by neurons with higher connectivities. The next example demonstrates this possibility.

Example 7. Consider $\eta = (+, +, +, +, +, +, -, -, -)$ ($p = 9$), and

$$\Sigma = (\eta^T, (\eta\mathbf{P})^T, (\eta\mathbf{P}^2)^T, (\eta\mathbf{P}^4)^T, (\eta\mathbf{P}^5)^T, (\eta\mathbf{P}^6)^T, (\eta\mathbf{P}^8)^T)^T.$$

This cycle is minimal as $\text{rank}(\eta) = \text{rank}(\Sigma) = 7$, but not consecutive. The gaps are between the third and fourth rows, and the sixth and seventh rows. Since the seventh and first rows are consecutive, there are no other gaps. The connectivity matrix is

$$\mathbf{J} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 1 & -1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and shows that we still have the forward chain 3 → 2 → 1 → 7 → 6 → 5 → 4, but neurons 3 and 6 receive multiple inputs. The network topology is shown in Figure 2C.

5.1.2. N-p relations

Definition 12. Let $N_p : \mathbb{X}^p \to \Omega$ be the function defined by $N_p(\eta) = n$ if η annihilates $(p-n)$ columns of V, where V is defined as in Theorem 1, $\mathbb{X}^p$ is the set of binary row vectors of length p, and $\Omega = \{1, 2, \dots, p-1\}$.

Remark 9. It is a direct consequence of Theorem 1 that $N_p(\eta) = \text{rank}(\eta)$. Therefore, for a given value of p, the image-set $N_p(\mathbb{X}^p)$ contains all possible values of N for which there exists $\eta \in \mathbb{X}^p$ such that (30) defines a simple MC-cycle. Furthermore, in Section 4.3 we have associated with $\eta \in \mathbb{X}^p$ the polynomial $p_\eta(x) = \eta(1, x, \dots, x^{p-1})^T$, where x is a complex variable. Since $p_\eta(\rho^k) = \eta v^{(k)}$, $\eta v^{(k)} = 0$ (i.e. η annihilates $v^{(k)}$) if and only if $p_\eta(x)$ has a factor which is a multiple of the minimal polynomial of ρ^k. Thus $N_p(\eta) = \text{rank}(\eta)$ is intimately related to the factorization of $p_\eta(x)$.

There appears to be no general characterization of or formula for $N_p(\eta)$. Even for row vectors with repeating block structure such as $\eta = (\sigma, -\sigma)$ or $(\sigma, -\sigma, \sigma)$ ($\sigma \in \mathbb{X}^{p/2}$ or $\mathbb{X}^{p/3}$), the factorization of $p_\eta(x)$ does not reveal a formalizable pattern. We therefore just list the sets $N_p(\mathbb{X}^p)$ in Table 1 for $1 \leq p \leq 20$. Note that $N_q(\mathbb{X}^q) \subset N_p(\mathbb{X}^p)$ if q divides p, since $\text{rank}(\eta) = \text{rank}(\sigma)$ if $\eta = (\sigma, \sigma, \dots, \sigma)$ (p/q repetitions) and $\sigma \in \mathbb{X}^q$. We therefore include in Table 1 only those values $N \in N_p(\mathbb{X}^p)$, for which there exists a row vector $\eta \in \mathbb{X}^p$ with $N = \text{rank}(\eta)$, and η is NOT a repetition of some shorter vector σ. To illustrate how Table 1 was obtained, we compute $N_p(\eta)$ for a row vector with $p = 6$ in Example 8.

Remark 10. Vectors η of the form $\eta = (\sigma, \sigma, \dots) \in \mathbb{X}^p$ with $\sigma \in \mathbb{X}^q$ have minimal period $\leq q$ under cyclic permutations. The number of binary vectors of minimal period p is found by subtracting the number of all vectors with smaller minimal period from 2^p. An inclusion/exclusion argument shows that this number is given by

$$M_p = 2^p - \sum_{k=1}^{s} (-1)^{k-1} \sum_{1 \leq i_1 < \cdots < i_k \leq s} 2^{p/(p_{i_1} p_{i_2} \cdots p_{i_k})},$$

if $p_1, p_2, \dots, p_s$ are the distinct prime numbers occurring in the prime factorization of p ($M_p = 2^p - 2$ if p is prime). Accordingly, the number of maximal loops, i.e. loops with $|\mathcal{L}_\eta| = p$, is M_p/p.

Example 8. Let $\eta = (+, +, -, -, -, +)$. This vector has a repeating block structure, $\eta = (\sigma, -\sigma)$, where $\sigma = (+, +, -)$. For $p = 6$ the matrix V is given

Table 1: Values of $N \in N_p(\mathbb{X}^p)$ attained by some $\eta \in \mathbb{X}^p$ that is NOT of the form $(\sigma, \sigma, \ldots, \sigma)$ with $\sigma \in \mathbb{X}^q$, $q < p$, for $p \leq 20$.

p	N	p	N
1	1	11	11
2	1	12	6,7,8,9,10,11,12
3	3	13	13
4	2,4	14	7,13,14
5	5	15	11,13,15
6	3,5,6	16	8,10,11,12,13,14,15,16
7	7	17	17
8	4,6,7,8	18	7,9,11,12,13,14,15,16,17,18
9	7,9	19	19
10	5,9,10	20	10,12,13,14,15,16,17,18,19,20

by

$$V = (v^{(0)}, v^{(1)}, \ldots, v^{(5)}) = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \rho & \rho^2 & -1 & \rho^4 & \rho^5 \\ 1 & \rho^2 & \rho^4 & 1 & \rho^2 & \rho^4 \\ 1 & -1 & 1 & -1 & 1 & -1 \\ 1 & \rho^4 & \rho^2 & 1 & \rho^4 & \rho^2 \\ 1 & \rho^5 & \rho^4 & -1 & \rho^2 & \rho \end{pmatrix},$$

where $\rho = e^{2\pi \mathbf{i}/6}$. The polynomial $p_\eta(x)$ has the following factorization,

$$\begin{aligned} p_\eta(x) &= 1 + x - x^2 - x^3 - x^4 + x^5 \\ &= (1 - x^3)(1 + x - x^2) \\ &= (1 - x)(1 + x + x^2)(1 + x - x^2). \end{aligned}$$

Since $\Phi_1(x) = x - 1$ and $\Phi_3(x) = x^2 + x + 1$ are the first and the third cyclotomic polynomials, and $\rho^0 = 1$ is the primitive first root of unity and ρ^2 and ρ^4 are the primitive third roots of unity, it follows that η annihilates $v^{(0)}$, $v^{(2)}$ and $v^{(4)}$. Therefore, $N_6(\eta) = 6 - 3 = 3$.

Some of the N-values in $N_p(\mathbb{X}^p)$ in Table 1 can be explained directly, without factorizing $p_\eta(x)$. We summarize three simple but important facts.

Proposition 8. *(a)* $\{1, p\} \subset N_p(\mathbb{X}^p)$ *for any* $p > 2$.
(b) If $p > 2$ *is prime, then* $N_p(\mathbb{X}^p) = \{1, p\}$.
(c) $p - 1 \in N_p(\mathbb{X}^p)$ *if* p *is even and* $p > 4$.

Proof. Since $\operatorname{rank}(+,+,\dots,+) = 1$, it follows that $1 \in N_p(\mathbb{X}^p)$ for any p. To show that $p \in N_p(\mathbb{X}^p)$ for $p > 2$, consider $\eta = (-,+,+,\dots,+)$ and let Σ_η be the $p \times p$-matrix defined by $\operatorname{row}_j(\Sigma_\eta) = \eta(\mathbf{P}^T)^{j-1}$, $1 \le j \le p$. By induction, one shows that $\det(\Sigma_\eta) = (-2)^{p-1}(p-2)$ which completes the proof of (a).

Statement (b) is an immediate consequence of the fact that $\Phi_p(x) = \sum_{j=1}^{p} x^{j-1}$ is the minimal polynomial of ρ^k, $0 < k < p$, if p is prime and is irreducible over $\mathbb{Q}$, hence if $\eta \neq \pm(+,+,\dots,+)$, $\Phi_p(x)$ and $p_\eta(x)$ cannot contain a common factor.

To show (c), let $\sigma = (+,-,+,-,\dots,+,-) \in \mathbb{X}^{p-2}$ and set $\eta = (\sigma,-,+)$. By performing elementary row operations on the matrix Σ_η with rows $\operatorname{row}_i(\Sigma_\eta) = \eta\mathbf{P}^{i-1}$, $1 \le i \le p-1$, it can be shown that Σ_η has full rank. The details are straightforward but tedious to write down explicitly and will be omitted. $\square$

Remark 11. (a) If p is prime, then $\operatorname{rank}(\eta) = p$ for any $\eta \neq \pm(+,+,\dots,+)$. For non-prime values of p one also can construct several different vectors with $\operatorname{rank}(\eta) = p$. For example, if p is odd, then $\operatorname{rank}(\eta) = p$ if $\eta = (+,-,+,-,\dots,+,-,+)$, which is easily shown using elementary row operations. A generalization is provided by vectors η with $\sum_i \eta^i = 1$. All our case studies indicate that these vectors have $\operatorname{rank}(\eta) = p$ as well.
(b) For even $p > 4$, the vector constructed in the proof of Proposition 8(c) is just one example of a vector with $\operatorname{rank}(\eta) = p - 1$. In general, if $\eta = (\eta^1,\dots,\eta^p)$ and $\sum_i \eta^i = 0$, then η is orthogonal to $(+,+,\dots,+)$, and $\operatorname{rank}(\eta) \le p - 1$. Case studies indicate that such a vector has maximal rank $p - 1$ if it does not have a "repeating block structure".

5.1.3. Simple anti-symmetric cycles

The characteristics of the cycles considered by Gencic et al. (1990) are that the cycle length p is even and the second $p/2$ columns of the cycle matrix are the negatives of the first $p/2$ columns in the same order. We call such cycles *anti-symmetric.* Here we discuss the possible values of the rank of the cycle matrix if these cycles are simple and admissible.

Proposition 9. *Assume p is even, $p = 2n$, and $\eta = (\sigma, -\sigma)$ with $\sigma = (\sigma^1,\dots,\sigma^n) \in \mathbb{X}^n$. Then we have the following possibilities for* $\operatorname{rank}(\eta)$.



Figure 2: Topology of networks constructed, respectively, by a simple MC-cycle with $N = p/2 = 7$ (**A**, Example 6), a simple MC-cycle with $N = p - 1 = 5$ (**B**, Example 6), a minimal simple but non-consecutive cycle with $N = p - 2 = 7$ (**C**, Example 7), a separable MC-cycle (**D**, Example 10) and a minimal genuinely inseparable composite cycle (**E**, Example 11). In panels **A-D**, excitatory (inhibitory) synaptic connections are labeled with red (blue) lines with arrowhead indicating the direction of the connections. In panel **E**, in order to highlight the clusters in the network, connections within the same clusters are labeled with black lines, and connections between neurons in different clusters are labeled with dark red lines. Directions and polarities of the connections are not shown in **E**. For all of the 5 networks illustrated in this figure, self-connectivities are not included.

(a) $1 \leq \operatorname{rank}(\eta) \leq n$. *Moreover,* $d \equiv n - \operatorname{rank}(\eta)$ *is even, and if* $d \geq 2$ *the* σ^j *satisfy* d *linearly independent homogeneous linear equations with integer coefficients.*
(b) If $p = 2^k$, $k \geq 2$, *then* $\operatorname{rank}(\eta) = n$.
(c) If $p = 2n$ *with* $n > 2$ *prime and* $\sigma \neq \pm(+, -, +, \dots, +, -, +)$, *then* $\operatorname{rank}(\eta) = n$.

Proof. (a) Let $\eta = (\sigma, -\sigma)$, $\sigma \in \mathbb{X}^n$, and define the $p \times p$-matrix Σ by $\operatorname{row}_j(\Sigma) = \eta(\mathbf{P}^T)^{j-1}$, $1 \leq j \leq p$. This matrix is a circulant matrix and contains all rows of $\mathcal{L}_\eta$, hence $\operatorname{rank}(\eta) = \operatorname{rank}(\Sigma)$. According to the properties of circulant matrices, the eigenvalues of Σ are of the form

$$\lambda = \sum_{i=1}^{p} \eta^i \rho^{i-1} = \sum_{i=1}^{n} (\sigma^i - \sigma^i \rho^n) \rho^{i-1}, \tag{33}$$

where ρ is any pth root of unity. Thus $\operatorname{rank}(\eta)$ coincides with the number of distinct pth roots of unity for which the right-hand side of (33) is nonzero. Since $\eta\mathbf{P}^n = -\eta$, it follows that $\operatorname{rank}(\eta) \leq n$, and clearly $\operatorname{rank}(\eta) \geq 1$, which proves the first statement of (a). To complete the proof of (a), we note that the $2n$ distinct roots of $x^{2n} = 1$ ($x \in \mathbb{C}$) comprise n roots with $x^n = 1$ and n roots with $x^n = -1$. Thus $d = n - \operatorname{rank}(\eta)$ coincides with the number of distinct roots of $x^n = -1$ for which

$$p_\sigma(x) = \sum_{i=1}^{n} \sigma^i x^{i-1} = 0. \tag{34}$$

If n is odd, $p_\sigma(-1) \neq 0$, and if n is even, $(-1)^n = 1$, thus all roots in question have nonzero imaginary parts, which implies that d is even. If $d \geq 2$, $p_\sigma(x)$ is divisible by a cyclotomic polynomial $\Phi_m(x)$, where m divides n but not $2n$. The degree of $\Phi_m(x)$ is given by Euler's totient function, $\varphi(m)$, and is even. The condition that $p_\sigma(x)$ factors through $\Phi_m(x)$ then leads to $\varphi(m)$ linearly independent homogeneous equations that must be satisfied by the σ^i, and since $\Phi_m(x)$ has integer coefficients, the coefficients of these equations can be chosen as integers as well. If $p_\sigma(x)$ contains several cyclotomic polynomials $\Phi_{m_j}(x)$, $1 \leq j \leq r$, as factors, the number of linear equations satisfied by σ is $\varphi(m_1) + \cdots + \varphi(m_r)$, and all these relations are linearly independent as the cyclotomic polynomials are distinct and irreducible over the rationals.
(b) If $p = 2^k$, $k \geq 2$, the only factor that divides p but not $n = 2^{k-1}$ is p. The cyclotomic polynomial of p is $\Phi_p(x) = 1 + x^n$ and has degree n, that is, $\Phi_p(x)$ cannot be a factor of $p_\sigma(x)$ which has degree $n - 1$.

(c) Assume now that $n > 2$ is prime. In this case, $\Phi_{2n}(x) = \Phi_n(-x)$ is the only cyclotomic polynomial in question and is given by $\Phi_n(-x) = 1-x+x^2-x^3+\cdots+x^n$. Thus, if σ is not of the form $\sigma = \pm(+,-,+,-,\ldots,+,-,+)$, $\Phi_n(-x)$ does not factor through $p_\sigma(x)$. $\square$

Since for $\eta = (\sigma,-\sigma)$ a "rank deficiency" ($\text{rank}(\eta) < n$) occurs only if σ satisfies a system of linear equations, the number of σ's for which $\text{rank}(\eta) = n$ is considerably larger than the number of σ's for which η has a rank-deficiency. Thus "generically" we expect that vectors of the form $(\sigma,-\sigma)$ have full rank n. The vector $\sigma = (+,-,+,-,\ldots,+,-,+)$ is, of course, a very special case as $(\sigma,-\sigma)$ has the repeating block structure $(+,-,+,-,\ldots,+,-)$ which has minimal rank 1. We illustrate the occurrence of rank deficiencies by an example.

Example 9. Let $p = 18 = 2\cdot 3^2$, i.e. $n = 9$. The cyclotomic polynomials that can give rise to a rank deficiency are here $\Phi_6(x) = 1-x+x^2$ and $\Phi_{18}(x) = 1-x^3+x^6$. The condition that $p_\sigma(x) = \sigma^1+\sigma^2x+\cdots+\sigma^9x^8$ factors through $\Phi_6(x)$ leads to the equations

$$\begin{aligned}\sigma^1-\sigma^3-\sigma^4+\sigma^6+\sigma^7-\sigma^9 &= 0,\\ \sigma^2+\sigma^3-\sigma^5-\sigma^6+\sigma^8+\sigma^9 &= 0.\end{aligned}$$

The only binary vector satisfying these conditions (up to cyclic permutations) are $\pm\sigma^{(1)}$, $\pm\sigma^{(2)}$ and $\pm\sigma^{(3)}$, where

$$\begin{aligned}\sigma^{(1)} &= (+,+,+,+,+,+,+,-,+),\\ \sigma^{(2)} &= (+,+,+,-,+,+,-,-,+),\\ \sigma^{(3)} &= (+,+,-,+,+,-,+,-,+).\end{aligned}$$

Since $\Phi_6(x)$ has a single pair of complex conjugate roots, $\text{rank}(\sigma^{(\nu)},-\sigma^{(\nu)}) = 9-2 = 7$, $\nu = 1,2,3$. Similarly, in order that $\Phi_{18}(x)$ factors through $p_\sigma(x)$, the conditions $\sigma^j+\sigma^{j+3} = 0$ for $1 \le j \le 6$ must be satisfied, leading to $\text{rank}(\sigma,-\sigma) = 3$. All vectors with block structure $\sigma = (\tilde\sigma,-\tilde\sigma,\tilde\sigma)$ with $\tilde\sigma \in \mathbb{X}^3$ have this property, and lead to $\eta = (\hat\sigma,\hat\sigma,\hat\sigma)$ with $\hat\sigma = (\tilde\sigma,-\tilde\sigma)$, i.e. $\text{rank}(\eta) = \text{rank}(\hat\sigma)$. This includes $\tilde\sigma = (+,+,+)$ with $\text{rank}(\hat\sigma) = 3$, and $\tilde\sigma = (+,-,+)$ with $\text{rank}(\hat\sigma) = 1$. In the latter case, both $\Phi_6(x)$ and $\Phi_{18}(x)$ are factors of $p_\sigma(x)$.

The rank of generic vectors (without rank deficiency) of the form (σ,σ) or $(\sigma,-\sigma)$ is equal to the length of σ. The converse question is under which circumstances a vector $\eta \in \mathbb{X}^p$ with p even and $\text{rank}(\eta) = p/2$ has this form. We state two simple sufficient conditions for this property.

Proposition 10. *Assume $p = 2n$, $\eta \in \mathbb{X}^p$, and* $\operatorname{rank}(\eta) = n$. *If $p = 2^k$, $k \geq 2$, or $n > 2$ is prime, then η is either of the form (σ, σ) or $(\sigma, -\sigma)$ for some $\sigma \in \mathbb{X}^n$.*

Proof. Let $\eta = (\sigma, \hat{\sigma})$ with σ, $\hat{\sigma} \in \mathbb{X}^n$. We consider again the matrix Σ defined in the proof of Proposition 9 with eigenvalues

$$\lambda = \sum_{i=1}^{n} (\sigma^i + \rho^n \hat{\sigma}^i) \rho^{i-1},$$

where ρ is a pth root of unity, $p = 2n$. Assuming that $\operatorname{rank}(\eta) = n$, there exist precisely n distinct roots ρ of $x^{2n} = 1$ for which $\lambda = 0$. We decompose these roots again into roots satisfying $x^n = 1$ and $x^n = -1$, respectively, and set accordingly

$$\lambda_{\pm}(x) = \sum_{i=1}^{n} (\sigma^i \pm \hat{\sigma}^i) x^{i-1}.$$

(a) Assume that $p = 2^k$ ($n = 2^{k-1}$) for $k \geq 2$. If there exists a root ρ of $x^n = -1$ for which $\lambda = 0$, $\lambda(x)$ must contain the cyclotomic polynomial $\Phi_{2n}(x) = 1 + x^n$ as a factor, which is only possible if $\sigma^i - \hat{\sigma}^i = 0$ for all i, because $\lambda(x)$ has at most degree $n - 1$. Thus $\eta = (\sigma, \sigma)$ in this case. Conversely, assume that all roots ρ for which $\lambda = 0$ are roots of $x^n = 1$. Then $\lambda_+(x)$ must contain all cyclotomic polynomials $\Phi_\nu(x)$ for $\nu = 1, 2, \ldots, 2^{k-1}$ as factors. Since the product of these polynomials is $1 - x^n$, this cannot hold unless $\sigma^i + \hat{\sigma}^i = 0$ for all i, thus $\eta = (\sigma, -\sigma)$ in this case.
(b) The case $p = 2n$ with $n > 1$ prime is treated similarly. Here the cyclotomic polynomials which factor through $x^n - 1$ are $1 - x$ and $\Phi_n(x) = 1 + x + \cdots + x^{n-1}$, and the cyclotomic polynomials which factor through $1 + x^n$ are $1 + x$ and $\Phi_n(-x)$. Since $n > 1$, either $\Phi_n(x)$ is a factor of $\lambda_+(x)$ or $\Phi_n(-x)$ is a factor of $\lambda_-(x)$, which implies that either $\hat{\sigma} = \sigma$ or $\hat{\sigma} = -\sigma$. □

An extension of Proposition 10 to more general values of p appears highly nontrivial, because a multitude of cyclotomic polynomial have to be considered if the prime factorization of n is more complicated. We have examined all vectors η with $\operatorname{rank}(\eta) = p/2$ for $p \leq 20$ and found that all these vectors have the form (σ, σ) or $(\sigma, -\sigma)$. Other vectors with $\operatorname{rank}(\eta) = p/2$ may exist for larger values of p, but if so we expect the number of these vectors to be much smaller than the number of (σ, σ)- or $(\sigma, -\sigma)$-vectors of full rank.

5.2. Separable MC-Cycles

For separable MC-cycles with generators $G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\}$, the spaces $\mathrm{span}\{\mathcal{L}_{\eta_j}\}$ and $\mathrm{span}\{\mathcal{L}_{\eta_k}\}$ intersect trivially if $j \neq k$. If Σ is in standard form, this implies immediately that $\mathbf{J}$ has a block structure, $\mathbf{J} = \mathrm{diag}(\mathbf{J}_1, \ldots, \mathbf{J}_r)$, where $\mathbf{J}_k$ is an $N_k \times N_k$-matrix of the form (31) with $N_k = \mathrm{rank}(\eta_k)$. Accordingly, a network constructed from a separable MC-cycle is decomposed into r disconnected clusters and for each cluster the connectivity matrix has the form corresponding to a simple MC-cycle.

Example 10. Consider the 7 × 8-cycle

$$\Sigma = \begin{pmatrix} + & + & + & + & - & - & - & - \\ + & + & + & - & - & - & - & + \\ + & + & - & - & - & - & + & + \\ + & - & - & - & - & + & + & + \\ + & + & - & - & + & + & - & - \\ + & - & - & + & + & - & - & + \\ + & - & + & - & + & - & + & - \end{pmatrix}.$$

This cycle has the generators η_1, η_5, η_7 ($\eta_j = \mathrm{row}_j(\Sigma)$) and is separable and in standard form. Moreover, $\eta_4\mathbf{P} = -\eta_1$, $\eta_6\mathbf{P} = -\eta_5$, and $\eta_7\mathbf{P} = -\eta_7$. Thus the network is decomposed into three clusters consisting of neurons 1, 2, 3, 4, neurons 5, 6, and neuron 7, with cycle-connectivity matrices

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

and -1, respectively. The topology of this network is illustrated in Figure 2D. We note, however, that the cluster consisting of neuron 7 cannot show oscillations without delay, since a $1D$ dynamical system does not have limit cycles. By contrast, with delay included, we can find oscillations already for $N = 1$ for appropriate parameter values (Zhang et al., 2013).

General separable cycles still can be decomposed into isolated clusters as $\mathrm{span}\{\mathrm{R}(\Sigma)\}$ is semisimple, however, the network topology in each cluster maybe more complicated (see Subsection 5.4). The issue with separable cycles is that, even if the subcycles corresponding to the different clusters are retrieved, these oscillations are in general not synchronized. We comment on this issue further in Section 6.

5.3. Minimal Inseparable Cycles

For minimal inseparable cycles Σ with generators $G_\Sigma = \{\eta_1, \eta_2, \ldots, \eta_q\}$, at least two subspaces $\mathrm{span}\{\mathcal{L}_{\eta_j} \cap \mathrm{R}(\Sigma)\}$ and $\mathrm{span}\{\mathcal{L}_{\eta_k} \cap \mathrm{R}(\Sigma)\}$ $(j \neq k)$ have a nontrivial intersection. Accordingly, Σ does not have full row-rank,

$$\mathrm{rank}(\Sigma) < \sum_{i=1}^{q} \mathrm{rank}(\eta_i) = N,$$

which implies in particular that $\mathbf{J^0} \neq I$. It is still possible to partition the network into clusters, but some clusters may be connected and the network topology within the cluster corresponding to the loop $\mathcal{L}_{\eta_j}$ will in general not coincide with the network topology predicted by the submatrix Σ_j of the corresponding simple cycle. Thus the consecutiveness requirement does not have an effect, whereas the minimality requirement takes care that the sub-matrices in $\mathbf{J^0}$ and $\mathbf{J}$ defining the connectivities within the clusters are non-singular. The following example illustrates these features.

Example 11. Consider the 10×12-cycle $\Sigma = (\Sigma_1^T, \Sigma_2^T, \Sigma_3^T)^T$, where

$$\begin{aligned} \Sigma_1^T &= (\eta_1^T, (\eta_1\mathbf{P})^T, (\eta_1\mathbf{P}^2)^T, (\eta_1\mathbf{P}^3)^T)^T, \\ \Sigma_j^T &= (\eta_j^T, (\eta_j\mathbf{P})^T, (\eta_j\mathbf{P}^2)^T)^T,\ j = 2, 3, \end{aligned}$$

with

$$\begin{aligned} \eta_1 &= (\, + \;\; + \;\; + \;\; - \;\; + \;\; + \;\; + \;\; - \;\; + \;\; + \;\; + \;\; - \,), \\ \eta_2 &= (\, - \;\; + \;\; + \;\; + \;\; - \;\; - \;\; - \;\; + \;\; + \;\; + \;\; - \;\; - \,), \\ \eta_3 &= (\, + \;\; - \;\; - \;\; + \;\; - \;\; - \;\; + \;\; - \;\; - \;\; + \;\; - \;\; - \,). \end{aligned}$$

This is a minimal inseparable admissible cycle with generator $G_\Sigma = \{\eta_1, \eta_2, \eta_3\}$. The connectivity matrix $\mathbf{J}$ is given by

$$\mathbf{J} = \frac{1}{8}\begin{pmatrix} 0 & 7 & 0 & -1 & 1 & -1 & 1 & -1 & -1 & -1 \\ -1 & 0 & 7 & 0 & -1 & 1 & -1 & -1 & -1 & -1 \\ 0 & -1 & 0 & 7 & 1 & -1 & 1 & -1 & -1 & -1 \\ 7 & 0 & -1 & 0 & -1 & 1 & -1 & -1 & -1 & -1 \\ 1 & -1 & 1 & -1 & 2 & 6 & 2 & 0 & 0 & 0 \\ -1 & 1 & -1 & 1 & -2 & 2 & 6 & 0 & 0 & 0 \\ 1 & -1 & 1 & -1 & 6 & -2 & 2 & 0 & 0 & 0 \\ -1 & -1 & -1 & -1 & 0 & 0 & 0 & -2 & 6 & -2 \\ -1 & -1 & -1 & -1 & 0 & 0 & 0 & -2 & -2 & 6 \\ -1 & -1 & -1 & -1 & 0 & 0 & 0 & 6 & -2 & -2 \end{pmatrix},$$

and $\mathbf{J^0}$ has the same block structure as $\mathbf{J}$ (with self-feedbacks of all neurons). From the form of $\mathbf{J}$ (and $\mathbf{J^0}$) we infer that the cluster corresponding to η_1 is connected to the clusters corresponding to η_2 and η_3, while the latter two clusters are not directly connected. This connectivity structure is due to the fact that $\mathrm{span}\{\mathcal{L}_{\eta_1}\}$ intersects $\mathrm{span}\{\mathcal{L}_{\eta_2}\}$ and $\mathrm{span}\{\mathcal{L}_{\eta_3}\}$ in the one-dimensional spaces spanned by $(+,-,+,-,\dots,+,-)$ and $(+,+,+,\dots,+)$, respectively, whereas $\mathrm{span}\{\mathcal{L}_{\eta_2}\}$ and $\mathrm{span}\{\mathcal{L}_{\eta_3}\}$ intersect trivially. The network topology for this example is shown in Figure 2E. In general, two clusters corresponding to two generators $\eta, \eta' \in G_\Sigma$ are connected, if there exists a sequence $\eta = \eta_1, \eta_2, \dots, \eta_{s-1}, \eta_s = \eta'$ of generators such that $\mathrm{span}\{\mathcal{L}_{\eta_j}\}$ and $\mathrm{span}\{\mathcal{L}_{\eta_{j+1}}\}$ intersect nontrivially for $0 \le j < s$.

5.4. Further Examples

The examples in this subsection serve to illustrate the possible effects of fewer essential generators than generators. Consider a cycle Σ with $EG_\Sigma = \{\epsilon_1, \dots, \epsilon_r\}$. If $r < |G_\Sigma|$, the loop vectors of at least one generator are contained in the span of the loop vectors of another essential generator. Assuming $|\mathrm{R}(\Sigma)| = N$, this implies

$$\mathrm{rank}(\Sigma) \le \sum_{i=1}^{r} \mathrm{rank}(\epsilon_i) < N,$$

and we encounter again a rank-deficiency that will destroy special structures in the clusters corresponding to the essential generators.

Example 12. The 6×6-cycle

$$\Sigma = \begin{pmatrix} + & + & - & - & + & - \\ + & - & - & + & - & + \\ - & - & + & - & + & + \\ - & + & - & + & + & - \\ + & - & + & + & - & - \\ + & - & + & - & + & - \end{pmatrix},$$

has two generators, $\mathrm{row}_1(\Sigma)$ and $\mathrm{row}_6(\Sigma)$, but $\mathrm{row}_6(\Sigma) = \mathrm{row}_1(\Sigma)+\mathrm{row}_3(\Sigma)+\mathrm{row}_5(\Sigma)$, thus there is only one essential generator, $\epsilon = \mathrm{row}_1(\Sigma)$. Without the sixth row, Σ would be a simple MC-cycle with ring-topology. The presence of the sixth row destroys this structure, which is revealed in the following forms of $\mathbf{J^0}$ and $\mathbf{J}$,

$$\mathbf{J^0} = \frac{1}{4}\begin{pmatrix} 3 & 0 & -1 & 0 & -1 & 1 \\ 0 & 4 & 0 & 0 & 0 & 0 \\ -1 & 0 & 3 & 0 & -1 & 1 \\ 0 & 0 & 0 & 4 & 0 & 0 \\ -1 & 0 & -1 & 0 & 3 & 1 \\ 1 & 0 & 1 & 0 & 1 & 3 \end{pmatrix}, \ \mathbf{J} = \frac{1}{4}\begin{pmatrix} 0 & 4 & 0 & 0 & 0 & 0 \\ -1 & 0 & 3 & 0 & -1 & 1 \\ 0 & 0 & 0 & 4 & 0 & 0 \\ -1 & 0 & -1 & 0 & 3 & 1 \\ -1 & -4 & -1 & -4 & -1 & -3 \\ -1 & 0 & -1 & 0 & -1 & -3 \end{pmatrix}.$$

Example 13. The cycle

$$\Sigma = \begin{pmatrix} 1 & -1 & 1 \\ -1 & 1 & -1 \\ 1 & 1 & 1 \end{pmatrix},$$

has three generators and one essential generator that can be chosen as the first or second row. Since Σ is non-singular, Σ is admissible and $\mathbf{J^0}$ is the identity matrix. A successfully retrieved cycle shows three consecutive phases 1, 2, and 3 during an oscillation. In phases 1 and 2, neuron 1 is "on" (+) and in phase 3 it is "off" (−), while neuron 2 is "on" in phase 1 and "off" in phases 2 and 3. Clearly, neuron 3 is "on" during all 3 phases. The matrix $\mathbf{J}$ is given by

$$\mathbf{J} = \begin{pmatrix} -1 & 1 & 1 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and shows that neurons 1 and 2 form an excitatory/inhibitory pair, whereas neuron 3 acts excitatory on neuron 1. Without this third neuron the oscillations of neurons 1 and 2 as required by the first two rows of Σ could not be implemented, since the submatrix of Σ consisting of these rows is not admissible.

6. Discussion and Conclusion

In this paper we have studied the structural features of admissible cycles and their relation to the topology of the corresponding networks. Our study was pursued in the context of continuous-time Hopfield-type networks, but the results apply to the discrete networks considered by Personnaz et al.

(1986) and Guyon et al. (1988) as well, since the conditions on the connectivity matrix in these networks are the same as the conditions imposed on the cycle component of the connectivity matrix in the continuous case. In this paper we have restricted our study to single cycles; an extension to storing multiple cycles is given in (Zhang et al., 2013).

We have shown that if and only if the discrete Fourier transform $\hat{\Sigma} = \Sigma V$ of a cycle matrix Σ contains exactly r nonzero columns, where $r = \text{rank}(\Sigma)$, then a continuous-time asymmetric Hopfield-type network can be constructed with the pseudoinverse learning rule. Based on the structural analysis of the invariant subspaces of the row space of Σ, the admissible cycles have been classified into simple cycles, and separable and inseparable composite cycles. This classification was based on the decomposition of the row space of Σ into subsets corresponding to disjoint loops. The admissibility of a cycle implied that all vectors of a loop are in the row space of Σ if Σ contains some of these loop vectors. If no loop-space associated with Σ is a subspace of another loop-space (the generators are essential generators), we have identified for each loop the neurons associated with the loop vectors contained in Σ with a cluster. For general admissible cycles the clusters are connected, and the connectivity of the clusters depends on the intersections of their loop-spaces. Two clusters are directly connected if their indecomposable invariant subspaces intersect non-trivially. They are "indirectly" connected if they are part of a chain of directly connected clusters.

If an admissible cycle is separable, the clusters are completely isolated. In this case each cluster corresponds to a simple cycle associated with a generator of Σ. If the simple cycle is minimal and consecutive, the cluster has the form of a feedforward chain from the last neuron to the first neuron with feedbacks to the last neuron from the other neurons. If in addition the length of the cycle, p, is even and the rank of the generator is $p/2$, we generically find a ring structure with excitatory or inhibitory connection from the first neuron to the last neuron, but we cannot exclude that special loops with these properties exist for which no ring-structure occurs. If the simple cycle is minimal but non-consecutive, we find more than one feedforward chains. Regarding non-minimal simple as well as composite cycles, it would be interesting to find equivalence relations similar to those of Golubitsky et al. (2005), and Golubitsky & Stewart (2006), relating networks constructed from non-minimal cycles to networks constructed from minimal cycles.

The complete isolation of the clusters of separable cycles means that each cluster has its own subcycle. The issue is that we cannot expect the different subcycles to synchronize, preventing the network to traverse the cycle states in the order prescribed by the cycle matrix. In this case an additional synchronization mechanism must be introduced to enforce synchrony. Such a mechanism can be in the form of a small coupling among the clusters or through an external periodic input acting as pacemaker, see Zhang et al. (2013).

The generation of cyclic patterns in biological neural networks is associated with central pattern generators (CPGs), and the storage and retrieval of cyclic patterns are fundamentally important in animal neural systems.

In animal neural systems, the CPG networks with different topologies have been experimentally identified in different animal species (e.g., MacKay-Lyons, 2002). Moreover, as some animal movements, such as swallowing, gastrointestinal motility etc., often require the coordination of several functional groups of muscles, some CPG networks controlling these muscles can reorganize during different phases of the movement (Dickinson & Moulins, 1992; Meyrand et al., 1994). Such CPG networks consist of pools of neurons that can function in several CPGs involved in the organization of various motor behaviors.

Our study showed that the topology of networks constructed from an admissible cycle is determined by structural features (decomposition into invariant indecomposable subspaces) of the cycle matrix. Networks constructed from cycles with different structural features have different network topologies. If animal movements change, the structural features of the corresponding cyclic patterns change as well, which requires reorganization of the CPGs. The analysis of these structural features and their implications on the network topology may provide a scenario for studying the reorganization of CPGs during different phases of movements, when the dynamics of single neurons does not play a significant role in generating cycle patterns.

Appendix A. Cyclotomic Polynomials

We summarize here the basic properties of the cyclotomic polynomials used in Sections 3-5, for details see Dummit & Foote (2004).

The cyclotomic polynomial of order p is defined by $\Phi_p(x) = \prod_r (x - x_r)$, $x \in \mathbb{C}$, where the x_r encompass all primitive p-th roots of unity, that is, $x_r^p = 1$ and $x_r^n \neq 1$ if $1 \leq n < p$. The total number of such primitive roots is given by *Euler's totient function*, $\varphi(p)$. If $p = \prod_j p_j^{m_j}$ with distinct primes p_j is the prime factorization of p, then $\varphi(p) = \prod_j p_j^{m_j - 1}(p_j - 1)$. The important property of the cyclotomic polynomial is that they have integer coefficients and are irreducible over the rationals. Moreover, $\Phi_p(x)$ is the minimal polynomial for each root x_r, and the product of all $\Phi_d(x)$ for which d is a factor of p and $1 \leq d \leq p$ is $x^p - 1$. The only cyclotomic polynomials of odd degree are $\Phi_1(x) = 1 - x$ and $\Phi_2(x) = 1 + x$, all $\Phi_p(x)$ for $p > 2$ have even degrees as their primitive roots are all complex. Some basic properties of $\Phi_p(x)$ are:

$$\Phi_p(x) = \sum_{i=1}^{p} x^{i-1} \text{ if } p \text{ is prime,}$$

$$\Phi_{2p}(x) = \Phi_p(-x) \text{ if } p \text{ is odd,}$$

$$\Phi_p(x) = \Phi_q(x^{p/q}),$$

where q is the radical of p, i.e. the product of all distinct prime numbers occurring in the prime factorization of p. The last property implies in particular $\Phi_p(x) = 1 + x^n$ if $p = 2n = 2^k$, $k \geq 1$. The third cyclotomic polynomial is $\Phi_3(x) = 1 + x + x^2$ and has the roots $e^{\pm 2\pi \mathbf{i}/3}$.

References

Dickinson, P.S. & Moulins, M. (1992) Interactions and combinations between different networks in the stomatogastric nervous system. In: *Dynamic biological networks: the stomatogastric nervous system* Harris-Warrick, R.M., Marder, E., Selverston, A.I., and Moulins, M. (eds) Cambridge, MA: MIT, 139-160

Diederich, S. & Opper, M. (1987) Learning of correlated patterns in spin-glass networks by local learning rules. *Phys. Rev. Lett. 58*, 949-952

Dummit, D.S. & Foote, R.M. (2004) *Abstract Algebra* (3rd ed.) John Willey & Sons, Inc.: Hoboken, NJ

Gencic, T., Lappe, M., Dangelmayr, G. & Guettinger, W. (1990) Storing cycles in analog neural networks. In *Parallel Processing in Neural Systems and Computers*, Eckmiller, R., Hartmann, G. and Hauske, G. (eds.), North Holland, 445-450

Golubitsky, M., Stewart, I. & Török, A. (2005) Patterns of synchrony in coupled cell networks with multiple arrows. *SIAM J. Appl. Dynam. Sys. 4(1)*, 78-100

Golubitsky, M. & Stewart, I. (2006) Nonlinear dynamics of networks: the groupoid formalism. *Bull. Amer. Math. Soc., 43(3)*, 305-364

Guyon, I., Personnaz, L., Nadal, J.P. & Dreyfus, G. (1988) Storage and retrieval of complex sequences in neural networks. *Physical Review A, 38(12)*, 6365-6372

Hopfield, J.J. (1982) Neural networks and physical systems with emergent collective computational abilities. *Proc. Natl. Acad. Sci. USA 79*, 2554-2558

Hopfield, J.J. (1984) Neurons with graded response have collective computational properties like those of two-state neurons. *Proc. Natl. Acad. Sci. USA 81*, 3088-3092

Kanter, I. & Sompolinsky, H. (1987) Associative recall of memory without errors. *Phys. Rev. A 35*, 380-392

Kleinfeld, D. (1986) Sequential state generation by model neural networks. *Proc. Natl. Acad. Sci. USA 83*, 9469-9473

Kleinfeld, D. & Sompolinsky, H. (1988) Associative neural network model for the generation of temporal patterns, theory and application to central pattern generators. *Biophys. J. 54*, 1039-1051

Kohonen, T. (1988) *Self-organisation and associative memory.* (3rd ed.) Springer-Verlag: New York

Korn, H. & Faure, P. (2003) Is there chaos in the brain? II. Experimental evidence and related models. *Comptes Rendus Biologies 326(9)*, 787-840

Lancaster, P. & Tismenetsky, M. (1985) *The theory of matrices.* (2nd ed.) Academic Press: San Diego

Little, W.A. (1974) The existence of persistent states in the brain. *Math. Biosci. 19*, 101-120

López-Rodríguez, D., Mérida-Casermeiro, E. & Ortiz-de-Lazcano-Lobato, J.M. (2005) Hopfield network as associative memory with multiple reference points. *World Academy of Science, Engineering and Technology 7*, 62-67

MacKay-Lyons, M. (2002) Central pattern generation of locomotion: a review of the evidence. *Phys. Ther. 82*, 69-83

Marinaro, M., Scarpetta, S. & Yoshioka, M. (2007) Learning of oscillatory correlated patterns in a cortical network of a STDP-based learning rule. *Math. Biosci. 207*, 322-335

McEliece, R.J., Posner, E.C., Rodemich, E.R. & Venkatesh, S.S. (1987) The capacity of the Hopfield associative memory. *IEEE Transactions on Information Theory IT-33(4)*, 461-482

Meyrand, P., Simmers, J., & Moulins, M. (1994) Dynamic construction of a neural network from multiple pattern generators in the lobster stomatogastric nervous system. *J. Neurosci. 14*, 630-644

Miller, W. Jr. (1972) *Symmetry groups and their applications.* Academic Press: New York

Personnaz, L., Guyon, I. & Dreyfus, G. (1986) Collective computational properties of neural networks: new learning mechanisms. *Physical Review A, 34(5)*, 4217-4228

Rojas, R. (1996) *Neural networks - a systematic introduction.* Springer-Verlag: New York

Šíma, J. & Orponen, P. (2003) Continuous-Time symmetric Hopfield nets are computationally universal. *Neural Computation 15*, 693-733

Zhang, C., Dangelmayr, G. & Oprea, I. (2013) Storing Cycles in Hopfield-type Networks with Pseudoinverse Learning Rule - Bifurcation Analysis. In preparation

Zifan, A., Gharibzadeh, S. & Moradi M.H. (2007) Could dynamic attractors explain associative prosopagnosia? *Medical Hypotheses 68*, 1399-1405